

# ORTHOPAEDICS



FOR PRIMARY  
HEALTH CARE



**LION**

LEARNING INNOVATION VIA  
ORTHOPAEDIC NETWORKS

UNIVERSITY OF CAPE TOWN'S **ORTHOPAEDIC DEPARTMENT**

Editor: Michael Held

# Bone and Joint Infections Basics

Authors: Michael Held, Maritz Laubscher

---

## Introduction

Bone and joint infections can be **acute** or **chronic**. Acute infections should be treated as orthopaedic emergency. Especially for children and immunocompromised patients infections, acute infections can lead to severe systemic illness and can have detrimental long term morbidity, if not treated urgently. **Osteomyelitis** (infection of bone) can be divided into acute, sub-acute and chronic. In subacute osteomyelitis infection last from one to several months after which chronicity begins. Chronic osteomyelitis is characterized by progressive bone destruction and new bone apposition.

## Pathophysiology

### Hematogenous

Compared to children, adult osteomyelitis is rarely hematogenous, i.e., originated or transported by blood. When it occurs, it usually affects the spine. It is caused by microorganisms that seed the bone in the event of bacteremia. The most common organism found in hematogenous spread is *Staphylococcus aureus*. Hematogenous infections are most common in children. The metaphysis is the most common site because it is rich in blood supply (although this has sluggish flow) and is an actively growing part of bone. It also has relatively fewer phagocytes than the physis or diaphysis.

### Contiguous spread

This mode of contamination can be associated with previous surgery, an old non healing wound (diabetic foot, neuropathic ulcer) or a previous trauma. Infection spread by contiguity from adjacent tissue to bone.

### Direct inoculation

Open fractures, penetrating injuries or bone surgeries can be a direct source of infection resulting in osteomyelitis

## Risk factors

- Recent trauma or surgery
- Immunocompromised patients
- IV drug use
- Poor vascular supply
- Systemic conditions such as diabetes and sickle cell anaemia
- Peripheral neuropathy

## Classification

Osteomyelitis may be classified based on the duration of illness (acute versus chronic) and the mechanism of infection (haematogenous versus non-haematogenous).

and account for 80-90% of positive cultures. The table below shows common organisms of different age groups and patient populations with antibiotic choice, but this should be guided by local authorities.

## Bacteriology and Antibiotics

The most common infecting organism is staphylococcus aureus in all age groups

Patient cohort and bacteriology		Antibiotics
Neonates	S. aureus (MRSA), Group B strep Gram negatives	Cloxacillin (Fusidic Acid) 3rd gen cephalosporin
6 months - 4 years	S. aureus, K. kingae, H. influenzae (rare)	Cloxacillin + Ampicillin/3rd gen cephalosporin
> 4 years	S. aureus	Cloxacillin
Adults (acute)	S. aureus	
Penicillin allergy		Clindamycin
immunocompromised	S. aureus, S. pneumoniae, pseudomonas, fungal	Cloxacillin + 3rd gen cephalosporin Cloxacillin + 3rd gen cephalosporin Cefazolin is an alternative to Cloxacillin (if not available)
Sickle cell disease	S. aureus, Salmonella	
contiguous chronic osteomyeliti	The most common causative organism in s are: Enterobacteriaceae sp. Staphylococcus sp. Pseudomonas aeruginosa Enterococcus sp. In 45% of infections multiple organisms cultured	
heamatogenous chronic osteomyelitis	Staphylocossus aureua is the most common organism involved in adults (60-90%)	

Common organisms of different age groups and patient populations with antibiotic choice,

**Editor:** Michael Held

**Conceptualisation:** Maritz Laubscher & Robert Dunn - **Cover design:** Carlene Venter Creative Waves - **Developmental editing and design:** Vela and Phinda Njisane

## About the book

Informed by experts: Most patients with orthopaedic pathology in low to middle-income countries are treated by non-specialists. This book was based on a modified Delphi consensus study with experts from Africa, Europe, and North America to provide guidance to these health care workers. Knowledge topics, skills, and cases concerning orthopaedic trauma and infection were prioritized. Acute primary care for fractures and dislocations ranked high. Furthermore, the diagnosis and the treatment of conditions not requiring specialist referral were prioritized.

**The LION:** The Learning Innovation via orthopaedic Network (LION) aims to improve learning and teaching in orthopaedics in Southern Africa and around the world. These authors have contributed the individual chapters and are mostly orthopaedic surgeons and trainees in Southern Africa who have experience with local orthopaedic pathology and treatment modalities but also in medical education of undergraduate students and primary care physicians. To centre this book around our students, iterative rounds of revising and updating the individual chapters are ongoing, to eliminate expert blind spots and create transformation of knowledge.

Reference: Held et al. Topics, Skills, and Cases for an Undergraduate Musculoskeletal Curriculum in Southern Africa: A Consensus from Local and International Experts. JBJS. 2020 Feb 5;102(3):e10.

## Disclaimers

Although the authors, editor and publisher of this book have made every effort to ensure that the information provided was correct at press time, they do not assume and hereby disclaim any liability to any party for any loss, damage, or disruption caused by errors or omissions, whether such errors or omissions result from negligence, accident, or any other cause.

This book is not intended as a substitute for the medical advice of physicians. The reader should regularly consult a physician in matters relating to his/her health and particularly with respect to any symptoms that may require diagnosis or medical attention.

The information in this book is meant to supplement, not replace, Orthopaedic primary care training. The authors, editor and publisher advise readers to take full responsibility for their safety and know their limits. Before practicing the skills described in this book, be sure that your equipment is well maintained, and do not take risks beyond your level of experience, aptitude, training, and comfort level.

The individual authors of each chapter are responsible for consent and right to use and publish images in this book. The published work of this book falls under the [Creative Commons Attribution \(CC BY\) International 4.0 licence](#)

## Acknowledgements

Michelle Willmers and Glenda Cox for their mentorship.