

ORTHOPAEDICS



FOR PRIMARY
HEALTH CARE



LION

LEARNING INNOVATION VIA
ORTHOPAEDIC NETWORKS

EDITOR: **MICHAEL HELD**

UNIVERSITY OF CAPE TOWN'S **ORTHOPAEDIC DEPARTMENT**

Hip fractures and dislocations

by Johan le Roux, Kim Laubscher & Michael Held

Learning objectives

1. Low-energy hip fractures are often associated with significant osteoporosis.
2. Patients with hip fractures need a comprehensive medical investigation and care.
3. Timely surgery and early mobilisation reduce complications and mortality in hip fractures.

Introduction

Fractures of the acetabulum and proximal femur are considered hip fractures. These injuries, and hip dislocations, are debilitating and must be treated urgently to allow the patient to regain mobilisation and avoid the morbidity and mortality associated with being bedridden.

Assessment

Clinical

Most patients with hip injuries have groin pain. In some patients with undisplaced or impacted fractures, mobilisation is still possible. Patients with displaced fractures or hip dislocations are usually not able to bear weight or raise a straight leg. The pull of the hip muscles will leave the leg in a particular position.

With fractures, the leg is held in external rotation, abduction. With posterior dislocation (90% of dislocations) the leg held in flexion, adduction, internal rotation. Most commonly, the leg is shorter than the contralateral side.

Imaging

AP and lateral X-ray of the hip and a full-length femur film are needed to exclude associated injuries. **Disruption of Shenton's line, loss of joint congruence, and a difference in femur head shape are indirect signs of an injury.**

In non-displaced cases, the fracture may be occult and not visible on X-rays. MRI is the gold standard for occult fractures.

If this cannot be arranged within 24 hours, a CT scan is the next best choice.

In hip dislocations, CT scans help visualise associated fractures of the acetabulum, femur head or neck, and exclude loose bodies in the joint.



X-ray pelvis (AP): Left hip fracture with disruption of the Shenton's line (yellow)

Relevant anatomy

The hip capsule originates from the acetabular margin and attaches on the greater and lesser trochanters and the intertrochanteric line. The femur neck is intracapsular, surrounded by synovial fluid, and not covered in periosteum, making the healing potential of these **femoral neck fractures** very poor. Another problem is the disruption of the medial circumflex femoral artery (the main blood supply to the femoral head), leading to avascular necrosis of the femoral head. The fractures can be grouped into *displaced or undisplaced* fractures, which guide surgical treatment.

Peritrochanteric fractures occur between the lesser and greater trochanter. Therefore, they are extra-articular and do not cause medial circumflex femoral artery disruption and have a low risk of avascular necrosis.



AP right hip: Peritrochanteric fracture with an intact femur neck.

are usually high-energy injuries with other associated injuries. A structured ATLS approach should be used during examination and initial treatment. Urgent closed reduction in line with the deformity under adequate analgesia and sedation are crucial. Neurovascular examination and documentation before and after reduction are mandatory. If unable to achieve closed reduction, closed reduction under general anaesthesia and muscle relaxant in theatre is the next step. If still unsuccessful, open reduction must be performed.

Femoral head fracture

These are rare fractures and are usually associated with hip dislocations. The hip dislocation must be reduced initially. The fracture can be treated conservatively if it is in a non-weight bearing area, non-displaced and not associated with other injuries. Otherwise, fixation is required.



AP X-ray pelvis: Left hip dislocation. The Shenton's line is disrupted. The hip is internally rotated (lesser trochanter is not visible), and the femoral head appears smaller than the contralateral side (closer to the X-ray film).

Hip dislocation treatment

These

Hip fracture treatment

After excluding other injuries (especially distal radius fractures and head injuries) and after neuromuscular structures are assessed, the patient should be made comfortable and evaluated for a general anaesthetic (see table). The race against time starts now as early mobilisation leads to improved outcomes if patients are treated within 48 hours of injury, thus preventing life-threatening complications such as DVT/PE, urinary tract infections and pneumonia. This is considered a surgical emergency and most centres plan the operation on the **next available surgical list**. A continued, coordinated orthogeriatric and multidisciplinary management of these patients with investigations for osteoporosis and fall prevention remains the gold standard, which few receive.

Operative treatment

Non-operative treatment is only considered in patients who are unfit for anaesthesia and surgery.

For **neck of femur fractures**, open reduction and internal fixation are indicated for most physiologically young patients. Undisplaced fractures are treated with cannulated screws or a sliding hip screw. Arthroplasty is indicated for elderly patients with displaced fractures.

For **peri-trochanteric fractures**, a dynamic hip screw is done in stable fracture patterns. In unstable fractures, a cephalomedullary nail is indicated.

(oblique fractures which would displace with dynamic compression of the hip screw, fractures with subtrochanteric extension or medial comminution)

Investigations for patients with hip fractures
Skin traction with low weight (2kg)
Analgesia
Rehydration
Monitor urine output
Chest X-ray
Full blood count
Arterial blood gas (with respiratory compromise)
Echocardiogram
Cross-match



AP pelvic X-rays: Total hip replacement (above) and dynamic hip screw (below).

Editor: Michael Held

Conceptualisation: Maritz Laubscher & Robert Dunn

Cover design: Carlene Venter
(Creative Waves Brand Design)

Developmental editing and design:
Vela Njisane and Phinda Njisane

ABOUT THE BOOK

Informed by experts: Most patients with orthopaedic pathology in low to middle-income countries are treated by non-specialists. This book was based on a modified Delphi consensus study* with experts from Africa, Europe, and North America to provide guidance to these health care workers. Knowledge topics, skills, and cases concerning orthopaedic trauma and infection were prioritised. Acute primary care for fractures and dislocations ranked high.

Furthermore, the diagnosis and the treatment of conditions not requiring specialist referral were prioritised.

* Held et al. Topics, Skills, and Cases for an Undergraduate Musculoskeletal Curriculum in Southern Africa: A Consensus from Local and International Experts. JBJS. 2020 Feb 5;102(3):e10.

THE LION

The Learning Innovation via Orthopaedic Network (LION) aims to improve learning and teaching in orthopaedics in Southern Africa and around the world. These authors have contributed the individual chapters and are mostly orthopaedic surgeons and trainees in Southern Africa who have experience with local orthopaedic pathology and treatment modalities but also in medical education of undergraduate students and primary care physicians. To centre this book around our students, iterative rounds of revising and updating the individual chapters are ongoing, to eliminate expert blind spots and create transformation of knowledge.

DISCLAIMERS

Although the authors, editor and publisher of this book have made every effort to ensure that the information provided was correct at press time, they do not assume and hereby disclaim any liability to any party for any loss, damage, or disruption caused by errors or omissions, whether such errors or omissions result from negligence, accident, or any other cause.

This textbook is not intended as a substitute for the medical advice of physicians. The reader should regularly consult a physician in matters relating to his/her health and particularly with respect to any symptoms that may require diagnosis or medical attention.

The information in this book is meant to supplement, not replace, Orthopaedic primary care training. The authors, editor and publisher advise readers to take full responsibility for their safety and know their limits. Before practicing the skills described in this book, be sure that your equipment is well maintained, and do not take risks beyond your level of experience, aptitude, training, and comfort level.

The individual authors of each chapter are responsible for consent and rights to use and publish images in this book.

© The Authors 2021



Licensed under a Creative Commons Attribution (CC BY) 4.0 International Licence.

ACKNOWLEDGEMENTS

This work was carried out with the aid of a grant from the International Development Research Centre, Ottawa, Canada. Thanks to Johan Fagan, Michelle Willmers and Glenda Cox for their mentorship and support.