

ORTHOPAEDICS



FOR PRIMARY
HEALTH CARE



LION

LEARNING INNOVATION VIA
ORTHOPAEDIC NETWORKS

EDITOR: **MICHAEL HELD**

UNIVERSITY OF CAPE TOWN'S **ORTHOPAEDIC DEPARTMENT**

Knee dislocations

by Sithombo Maqungo, Abdalslam Andisha & Mlekeleli Duma

Learning objectives

1. Describe the clinical findings associated with knee dislocation.
2. Perform a closed reduction of knee dislocation effectively.

Introduction

Knee dislocations are traumatic injuries associated with a high rate of vascular injury. Treatment is generally emergent reduction and assessment of limb perfusion. However, there is a subset of patients, especially those with a high BMI, who can also sustain knee dislocations from low energy falls and a high index of suspicion is required in this regard. Often these injuries will lead to ligament laxity and stiffness.

Clinical assessment

The patient presents with knee pain, joint deformity, swelling and effusion.

Patients who sustain a knee dislocation that spontaneously reduces will show subtle signs of the injury such as abrasions and effusion. A suspicion of a patella dislocation or subluxation should be kept in mind.


A **vascular examination** needs to be performed to rule out popliteal artery injury. *Dorsalis pedis* and posterior tibial artery pulses should be evaluated.

If the pulse is absent, perform immediate closed reduction as this represents a threatened limb. A reduction should be performed before radiography. If the pulse is still absent after reduction, inform the vascular surgeon and go to the operating room for exploration. The orthopaedic surgeon will also be involved to assist with limb stabilisation, often using an external fixator.

Angiographic assessment is mandatory in centres where available. A sequential assessment of the Ankle Brachial Index (ABI) is needed in all cases with a pulse present. This is the ratio of the blood pressure measured at the level of the ankle or the arm. The normal value is above 0.9. The vascular status of the limb should be evaluated and documented before and after the reduction. A neurologic examination should be part of the assessment; sensory and motor function of the peroneal and tibial nerves must be assessed.

Imaging

Obtain **AP and lateral X-ray views**. With reduced knees, irregular/asymmetric joint space is often an indicator of ligament damage. Associated fractures can also be excluded. (See X-ray).



Picture of X-ray: An anterior knee dislocation is seen on these images.

A **CT angiogram** is often performed to rule out popliteal artery injury (See CT scan). Renal function should be optimised before injection of the contrast.



Picture of CT scan: Injected arterial contrast shows obliteration of the popliteal artery.

An **MRI** is invaluable in knee dislocations as the soft tissue injuries are better identified, which aids surgical management


planning. Injuries to the periarticular/ intra-articular ligamentous structures (cruciate and collateral ligaments, posteromedial and postero-lateral corner structures) and chondral and meniscal injuries are better identified. (See MRI scan)



Reduction

Traction and gentle manipulation of anterior or posterior translation is the best method to reduce a knee dislocation.

Posteromedial dislocation can be associated with buttonholing of medial femoral condyle through the joint capsule. This makes closed reduction very problematic, and open reduction in the operating room is often needed. The knee should be splinted at twenty degrees of flexion to maintain the position after reduction.



Schematic: Reduction techniques for knee dislocations.

Essential take-aways

A dislocated knee should be treated as an emergency as it represents a threatened limb.

Immediate reduction should be undertaken without waiting for radiography.

Popliteal artery disruption is seen in 20% to 60% of cases.

A dislocated knee may be part of multiple injuries in a patient and, therefore, ATLS principles should form part of the initial assessment and management of the patient.

Knee dislocations are disabling injuries, and they need appropriate early care to reduce morbidity.

Editor: Michael Held

Conceptualisation: Maritz Laubscher & Robert Dunn

Cover design: Carlene Venter
(Creative Waves Brand Design)

Developmental editing and design:
Vela Njisane and Phinda Njisane

ABOUT THE BOOK

Informed by experts: Most patients with orthopaedic pathology in low to middle-income countries are treated by non-specialists. This book was based on a modified Delphi consensus study* with experts from Africa, Europe, and North America to provide guidance to these health care workers. Knowledge topics, skills, and cases concerning orthopaedic trauma and infection were prioritised. Acute primary care for fractures and dislocations ranked high.

Furthermore, the diagnosis and the treatment of conditions not requiring specialist referral were prioritised.

* Held et al. Topics, Skills, and Cases for an Undergraduate Musculoskeletal Curriculum in Southern Africa: A Consensus from Local and International Experts. JBJS. 2020 Feb 5;102(3):e10.

THE LION

The Learning Innovation via Orthopaedic Network (LION) aims to improve learning and teaching in orthopaedics in Southern Africa and around the world. These authors have contributed the individual chapters and are mostly orthopaedic surgeons and trainees in Southern Africa who have experience with local orthopaedic pathology and treatment modalities but also in medical education of undergraduate students and primary care physicians. To centre this book around our students, iterative rounds of revising and updating the individual chapters are ongoing, to eliminate expert blind spots and create transformation of knowledge.

DISCLAIMERS

Although the authors, editor and publisher of this book have made every effort to ensure that the information provided was correct at press time, they do not assume and hereby disclaim any liability to any party for any loss, damage, or disruption caused by errors or omissions, whether such errors or omissions result from negligence, accident, or any other cause.

This textbook is not intended as a substitute for the medical advice of physicians. The reader should regularly consult a physician in matters relating to his/her health and particularly with respect to any symptoms that may require diagnosis or medical attention.

The information in this book is meant to supplement, not replace, Orthopaedic primary care training. The authors, editor and publisher advise readers to take full responsibility for their safety and know their limits. Before practicing the skills described in this book, be sure that your equipment is well maintained, and do not take risks beyond your level of experience, aptitude, training, and comfort level.

The individual authors of each chapter are responsible for consent and rights to use and publish images in this book.

© The Authors 2021



Licensed under a Creative Commons Attribution (CC BY) 4.0 International Licence.

ACKNOWLEDGEMENTS

This work was carried out with the aid of a grant from the International Development Research Centre, Ottawa, Canada. Thanks to Johan Fagan, Michelle Willmers and Glenda Cox for their mentorship and support.