



CLOSURE OF PAEDIATRIC TRACHEOCUTANEOUS FISTULA: SURGICAL TECHNIQUE

Buhlebenkosi Hlomani, Shazia Peer

A tracheocutaneous fistula (TCF) is an epithelial-lined communication between skin and trachea that persists after decannulation (removal) of a tracheostomy tube. TCF rates are higher in paediatric patients. It is a recognised complication of long-standing tracheostomies and has been linked to longer duration of tracheostomy dependence ¹.

Complications associated with a persistent TCF include skin irritation; inadequate glottic closure causing a weak cough and airway compromise; poor phonation; poor cosmesis; increased risk of aspiration during swimming, bathing *etc.* ². Addressing a persistent TCF avoids delays with integration into society and mainstream schooling.

TCF is due to squamous epithelialisation of the tracheostomy tract. Closure is achieved by excising the fistula followed either by primary closure or allowing for healing to occur by secondary intention. Although primary closure provides immediate resolution of the fistula and better cosmesis, it may be associated with life-threatening complications ³. Allowing healing by secondary intention minimises such potential complications. It may however be perceived as “inconvenient” for the patients.

It is the authors' preference to allow healing by secondary intention, safety in the immediate postoperative period being the primary reason.

Preoperative evaluation

1. Patient selection is as important as timing of surgery: We employ the following eligibility criteria based on aetiological and patient factors and use them as a guide for timing of surgery:

- The original reason for the upper airway obstruction that necessitated

the tracheostomy has resolved/been reversed

- The child has been successfully decannulated for >1 year; this sees the child through 1 winter season
- Typically, children are >4 years old, weigh >15kg and have no central neurological deficits or cardiovascular compromise *e.g.* post cardiac surgery or with pulmonary disease
- If the surgery is done on a 2 - 4-year-old child weighing >15kg, it is advisable to insert a tracheostomy tube immediately post-operatively for the 1st 24hrs
- Diagnostic assessment of the upper airways is routinely performed before any decision is taken about surgery. Anatomical and physiological factors are discussed later

2. Flexible nasendoscopy: This is typically done in the clinic with the child awake. The nasopharynx, oropharyngeal inlet, base of tongue (BOT), supraglottis and vocal cord mobility are carefully assessed.

3. Overnight sleep study / Overnight oximetry: This is useful to assess children with reported sleep disordered breathing. The TCF is covered while asleep to assess for proximal airway obstruction.

4. Direct laryngoscopy, tracheoscopy and bronchoscopy

- During induction of anaesthesia, the patient must breathe spontaneously and cope well with bag mask ventilation. The TCF can be covered with a dressing
- Flexible bronchoscopy is then done through a laryngeal mask
 - Spray the larynx with topical lignocaine

- Confirm cord mobility
- Perform dynamic assessment of upper airways, especially in cases of tracheomalacia or bronchomalacia
- When aspiration is a concern, a bronchoalveolar lavage (BAL) is done
- Perform rigid laryngoscopy +/- suspension
 - Assess all levels of the airway *i.e.* oropharynx, base of tongue, vallecula, epiglottis, supraglottis, false and true cords, subglottis, trachea (proximal to stoma), peristoma, and distal trachea to the carina
 - If pathology is identified, a staged approach with initial surgical correction may be required before the TCF can be addressed *e.g.* removing a suprastomal granuloma, addressing persistent subglottic stenosis *etc*
 - Record the size of the airway using appropriate-sized endotracheal tubes. A child of 4 years must admit at least a 4.5 ETT with a leak at 20cm H₂O. If older, use an uncuffed ETT size (Age/4) + 4.

If a child is considered eligible for closure of the TCF, then proceed to surgery

Surgical procedure to close a TCF

- Administer a single dose of prophylactic antibiotics (Cefazolin), and a single dose of intravenous steroids
- Intubate the airway with an appropriate-sized cuffed (0.5 size smaller to accommodate a cuff) north-facing endotracheal tube (*Figures 1ab*)
- Position the child supine with a shoulder roll to extend the neck (*Figures 1ab*)
- A head ring may be required to maintain the head in a neutral position

- Infiltrate the skin around the TCF with local anaesthetic (Lignocaine with adrenaline). Do not exceed the maximum safe dose (*Figure 2*)



Figure 1a: Child placed supine with neck extended and north facing ETT



Figure 1b: Patient with a prominent occiput positioned without a head ring



Figure 2: Infiltration with local anaesthetic

- After cleaning and draping, identify the midline structures as tracheostomy sites typically are in a paramedian position and a TCF may be further from the midline than expected (*Figure 3*)



Figure 3: The TCF may lie off the midline

- Mark a transverse ellipse of skin around the stoma (*Figure 4*)
- Incise the skin along the elliptical markings around the TCF
- Elevate the two ends and grip them using a tissue forceps (*Figure 5*)
- Dissect the soft tissue circumferentially around the stoma using cautery and sharp and blunt dissection
- Stay close to the TCF
- Expose a 10mm segment of trachea (*Figure 6*)
- Avoid dissecting into trachea
- Enter the trachea close to the TCF using an #11 blade, and circumferentially excise the TCF, removing as little tracheal tissue as possible (+/- 5mm) (*Figure 7*)



Figure 4: Mark the skin around the TCF as an ellipse



Figure 5: Skin incision, elevating the edges and grasping skin with tissue forceps



Figure 6: Excision of tract, down to trachea



Figure 7: Resected TCF

- Avoid a large defect by not excising too much of the anterior tracheal wall as it will take longer to close (*Figures 7, 8*)

- Ensure that there is no remaining skin in the tract as this will prevent closure
- Oppose the skin edges with *steristrips* (Figure 9)
- Cover the defect with a pliable dressing that allows for a reservoir of air when the child breathes (Figure 10)




Figure 8: Defect following resection of TCF



Figure 9: Approximate wound edges with *steristrips*



Figure 10: Cover the defect with a pliable dressing

Postoperative care

- The surgeon must accompany the child from the operating room to the recovery room with an unopened, appropriately sized paediatric tracheostomy tube that must be kept at the bedside
- Once fully recovered, the child must be nursed in a high care ward
- For the first 24 hours, continuous monitoring of oxygen saturation, respiratory rate and heart rate, and watching for signs of respiratory distress, are paramount
- Sedation and good analgesia are recommended as crying and coughing increase the risk of surgical emphysema
- If there is any breathing difficulty postoperatively, immediately remove the dressing and reinsert a tracheostomy tube
- The child is transferred to a general paediatric ward for 24-48 hours
- Give strict instructions to the caregiver to avoid water around the revised stoma
- Change the dressing if it becomes soiled
- If there are copious secretions it is preferable to keep the stoma uncovered
- The child is reviewed 1 week postoperatively...the stoma is typically healed by then (Figure 11)



Figure 11: Stoma healed by secondary intention 1-week postoperatively

Complications

Immediate / 1st 24hrs

- Respiratory distress
 - Pneumothorax (*Figure 12*)
 - Pneumomediastinum
- Surgical emphysema
 - Air escaping from trachea into surrounding tissues
 - May be forced into mediastinum and cause a pneumomediastinum
 - Immediate management includes prompt reinsertion of the tracheostomy tube and oxygen supplementation



Figure 12: Acute right-sided pneumothorax

Delayed

- Failure of TCF to close
- Peristomal granulation tissue
- Suprastomal (anterior) tracheal collapse
- Tracheomalacia
- Tracheal stenosis

References

1. Ha T, Goyal M, Ongkasuwan J. Duration of tracheostomy dependence and development of tracheocutaneous fistula in children. *The Laryngoscope*. 2017;127(12): 2709-12
2. Wine T, Simons JP, Mehta DK. Comparison of 2 Techniques of Tracheocutaneous Fistula Closure Analysis of Outcomes and Health Care Use. *JAMA Otolaryngol Head Neck Surg*. 2014;140(3):237-42
3. Osborn AJ, de Alarcón A, Hart CK, Cotton RT, Rutter MJ. Tracheocutaneous Fistula Closure in the Pediatric Population: Should Secondary Closure Be the Standard of Care? *Otolaryngology–Head and Neck Surgery*. 2013;149(5): 766–71

How to cite this chapter

Hlomani B, Peer S. (2019). Closure of paediatric tracheocutaneous fistula - surgical technique. In *The Open Access Atlas of Otolaryngology, Head & Neck Operative Surgery*. Retrieved from <https://vula.uct.ac.za/access/content/group/ba5fb1bd-be95-48e5-81be-586fbaeba29d/Closure%20of%20paediatric%20tracheocutaneous%20fistula%20-%20surgical%20technique.pdf>

Authors

Buhlebenkosi Hlomani MBChB
Paediatric Otolaryngology Fellow
Red Cross War Memorial Children's Hospital
Division of Otolaryngology
University of Cape Town
Cape Town, South Africa
b_hlomani@yahoo.com

Shazia Peer MBBCh, FCORL, MMed
Head, Paediatric Otolaryngology
Red Cross War Memorial Children's Hospital
Division of Otolaryngology
University of Cape Town
Cape Town, South Africa
shazia.peer@uct.ac.za

Editor

Johan Fagan MBChB, FCS (ORL), MMed
Emeritus Professor and Past Chair
Division of Otolaryngology
University of Cape Town
Cape Town, South Africa
johannes.fagan@uct.ac.za

**THE OPEN ACCESS ATLAS OF
OTOLARYNGOLOGY, HEAD &
NECK OPERATIVE SURGERY**
www.entdev.uct.ac.za



The Open Access Atlas of Otolaryngology, Head & Neck Operative Surgery by [Johan Fagan \(Editor\)](#) johannes.fagan@uct.ac.za is licensed under a [Creative Commons Attribution - Non-Commercial 3.0 Unported License](#)

