



Cricothyroidotomy, also known as cricothyrotomy refers to the creation of a communication between airway and skin via the cricothyroid membrane. It may be achieved by needle cricothyrotomy or by open or percutaneous cricothyroidotomy technique. Advantages of cricothyroidotomy compared to tracheostomy include simplicity, speed, relatively bloodless field, minimal training required, and avoiding hyperextending the neck in patients with possible cervical spinal injury.

Indications

1. Airway obstruction proximal to the subglottis
2. Respiratory failure
3. Pulmonary toilette in patients unable to clear copious secretions
4. Bronchoscopy

For indications (1) and (2), cricothyroidotomy is generally done as an **emergency temporising procedure** when a patient cannot be intubated, or when tracheostomy would be too time consuming or difficult. Following cricothyroidotomy the patient should be **intubated or a formal tracheostomy done within 24hrs** to avoid complications such as glottic and subglottic stenosis.

Contraindications

- Inability to identify surface landmarks (thyroid cartilage, cricoid, cricothyroid membrane) due to e.g. obesity, cervical trauma
- Airway obstruction distal to subglottis e.g. tracheal stenosis or transection
- Laryngeal cancer: Other than for an extreme airway emergency, avoid a

cricothyroidotomy so as not to seed the soft tissue of the neck with cancer cells

- Coagulopathy (other than emergency situation)

Cricothyroidotomy in children

The cutoff age beyond which surgical cricothyroidotomy can be safely performed is unclear. The most conservative cutoff age quoted is 12yrs; in young children the cricothyroid membrane is smaller, the larynx is more funnel-shaped, rostral, and compliant and cricothyroidotomy may be more prone to causing subglottic stenosis. Hence needle cricothyrotomy (12–14 gauge cannula passed over a needle) is preferred.

Surface anatomy (Figures 1a, b)

With the neck in a neutral or extended position, identify the midline thyroid prominence or “Adam’s apple”. Moving inferiorly, the next solid prominence in the midline is the cricoid cartilage. Immediately above the cricoid the finger slips into the depression of the cricothyroid membrane.




Figure 1a: Surface anatomy




Figure 1b: Surface anatomy

Surgical Anatomy

A cricothyroidotomy enters the larynx in the midline just below the vocal cords. The incision passes through skin, subcutaneous fat, middle cricothyroid ligament of cricothyroid membrane, and mucosa of subglottic larynx (Figure 2).




Figure 2: Relations of thyroid cartilage, cricothyroid membrane, cricoid cartilage,

and thyroid gland (Brown) to location of cricothyroidotomy (Yellow line)




Figure 3: Note proximity of tube to vocal cords, and relationship to cricoid ring

The tube then passes through the cricoid ring, which is the narrowest part of the upper airway (Figure 3, 4).




Figure 4: Note proximity of tube to vocal cords, and relationship to cricoid ring

The thyroid isthmus typically crosses the 2nd and 3rd tracheal rings, and is out of harm's way, unless a pyramidal thyroid lobe is present (*Figure 2*). The only blood vessels that may be encountered are the anterior jugular veins (off the midline), and the cricothyroid arteries.

The *cricothyroid artery* is a small branch of the superior thyroid artery and courses across the upper part of the cricothyroid membrane and communicates with the artery of the opposite side (*Figure 5*). **Therefore incise the membrane along the superior margin of the cricoid.**




Figure 5: Cricothyroid artery

The **dimensions of the cricothyroid membrane** have a bearing on the size of endotracheal or tracheostomy tube to be used; the outer diameter (OD) of the tube should not exceed the diameter of the cricothyroid opening so as to avoid injury to the larynx. Even though the cricothyroid membrane measures 30mm in the horizontal plane, the gap between the cricothyroid muscles through which the tube has to pass is much less (*Figure 6*). Based on studies of the dimensions of cricothyroid membranes^{1,2,3} it has been recommended that a tube of no more than 9-10mm OD be used¹; This corresponds to a 7mm ID tube. An alternative suggestion

is to select a tube which is 1mm smaller than would normally be used for oro-tracheal intubation². If a Shiley tracheostomy tube is to be used, it should not exceed Size 4 (9.4mm OD)




Figure 6: Dimensions of cricothyroid membrane: Range & mean values in millimetres¹

Needle cricothyrotomy

Needle cricothyrotomy using a **12 or 14-gauge cannula** (*Figure 7*) is only employed as an interim measure in an extreme emergency when one is unable to do an open cricothyroidotomy, and in children.



Figure 7: Example of intravenous cannula with needle removed from cannula

Ventilation may be effective if the cannula is attached to high pressure jet ventilation. Ventilation is controlled with

a hand-operated jet injector (Figure 8) attached to central wall oxygen, tank oxygen, or the fresh gas outlet of an anaesthesia machine; or may be controlled with the oxygen flush valve of an anaesthesia machine.



Figure 8: Hand-controlled high-pressure jet ventilator

However such ventilation via needle cricothyrotomy will only suffice for approximately 45mins as it does not permit adequate ventilation and hence leads to an accumulation of CO₂; this can be particularly deleterious for head injury patients as hypoventilation causes raised intracranial pressure. **Adult patients must therefore be either intubated or cricothyroidotomy or formal tracheostomy done within 45 minutes.**

Should jet ventilation not be available, one would attach either a ventilator or an ambubag to the cannula. **However ventilation with a low-pressure self-inflating resuscitation bag is ineffective within a minute or so.**

The cannula can be attached to an ambubag or ventilator in two simple ways:

1. Fit a 2 or 3ml syringe to the cannula with plunger removed; insert the connection piece of a 7.5mm ID endo-

tracheal tube into the barrel of the syringe (Figure 9)

2. Fit a 10ml plastic syringe to the cannula with plunger removed; insert an endotracheal tube into the barrel of the syringe and inflate the cuff (Figure 10)



Figure 9: Connection piece of 7.5mm endotracheal tube attached to barrel of 2 or 3ml syringe




Figure 10: Insert an endotracheal tube into the barrel of a syringe and inflate the cuff

Complications of needle cricothyrotomy include pneumothorax, subcutaneous and mediastinal emphysema, bleeding, oesophageal puncture, and respiratory acidosis due to hypoventilation. Complete upper airway obstruction proximal to the cricothyrotomy is a contraindication to needle cricothyrotomy because of the risk of causing barotrauma to the lungs. Long term complications include subglottic stenosis and vocal cord injury.

Needle cricothyrotomy: Surgical steps

1. Position the patient supine with neck exposed and extended (if possible)
2. Identify surface landmarks *i.e.* thyroid cartilage, cricoid cartilage and cricothyroid membrane

3. Prepare a sterile field
4. Inject 1% lidocaine with 1:100 000 epinephrine into the skin and through the cricothyroid membrane into the airway to anaesthetise the airway and suppress the cough reflex (if time to do so)
5. Fix the thyroid cartilage with the 1st & 3rd fingers of the non-dominant hand leaving the 2nd finger free to locate the cricothyroid membrane
6. With the dominant hand, pass a 14-gauge intravenous cannula attached to a syringe filled with normal saline, through the cricothyroid membrane, directing it caudally at 45^o (Figure 11). Bending the distal part of the needle can assist with directing the catheter along the tracheal lumen (Figure 12).



Figure 11: Fix the larynx and insert intravenous cannula at 45^o




Figure 12: Cannula has been bent for easier access

7. Apply negative pressure to the syringe as the needle is advanced. Air bubbles will appear in the fluid-filled syringe as the needle traverses the membrane and enters the trachea (Figure 13).




Figure 13: Air bubbles appear in the fluid-filled syringe as the needle traverses cricothyroid membrane⁴

8. Advance the cannula and then retract the needle
9. Attach jet ventilation and ventilate at 15 L/min
10. Judge the adequacy of ventilation by movement of the chest wall and auscultation for breath sounds, and by pulse oximetry

Open Surgical Cricothyroidotomy

Preoperative evaluation

- *Level of obstruction:* Cricothyroidotomy will not bypass obstruction in the trachea or bronchial tree
- *Coagulopathy:* Unless an emergency, a coagulopathy should be corrected prior to the procedure
- *Surface anatomy of the neck:* Are the relevant landmarks palpable?

Preoperative preparation

Prepacked cricothyroidotomy kits are available, both for patients requiring airway support (Figure 14) and for patients requiring access for suctioning excessive secretions (Figure 15). However in an emergency airway situation no. 11 or 15 surgical blades, knife handle, curved artery forceps and thin endotracheal tube will do.



Figure 14: Cricothyroidotomy kit for patients requiring airway support and ventilation: Small cuffed tracheostomy tube, syringe, scalpel, T-piece, lubricant gel, suture, tracheostomy tape




Figure 15: Mini-tracheostomy kit for patients only requiring lavage and suctioning of secretions: 4mm uncuffed tracheostomy tube with introducer, scalpel, tracheostomy tape, connector piece and suction tubing

Open surgical cricothyroidotomy: Surgical steps

1. Position the patient supine with anterior neck exposed and extended (if possible)
2. Identify the surface landmarks *i.e.* thyroid cartilage, cricoid cartilage and cricothyroid membrane
3. Prepare a sterile field
4. Inject 1% lidocaine with 1:100 000 epinephrine into the skin, soft tissue and through the cricothyroid membrane into the airway to anaesthetise the airway (if there time)
5. Fix the thyroid cartilage with the 1st and 3rd fingers of the non-dominant

hand leaving the 2nd finger free to palpate the cricothyroid membrane

6. If the surface anatomy is well defined use the dominant hand to make a 1-2cm transverse stab incision with a scalpel directly over and right through the cricothyroid membrane at the superior margin of the cricoid (*Figure 15*). In the more thickset patient, make a 3cm vertical midline incision extending inferiorly from the thyroid prominence (*Figure 16*); dissect bluntly down to the cricothyroid membrane with the non-dominant index finger; move the finger from side-to-side to clearly feel the cricothyroid membrane




Figure 16: Horizontal or vertical skin incision

7. Make a 1cm transverse incision through the cricothyroid membrane along the superior edge of the cricoid, angling the scalpel cephalad so as to avoid injuring the vocal cords; await a distinct “pop” sensation as the scalpel pierces through the membrane and enters the larynx; the thick cricoid lamina is located directly posteriorly (*Figures 3, 4*)
8. Dilate the tract by passing a curved haemostat through the incision, angling it caudad through the cricoid ring and along the trachea taking care not to perforate the posterior wall of the

trachea (Figure 4); alternatively insert a bougie through the tract into the airway

9. Insert a tracheostomy or endotracheal tube (≤ 7 mm ID), either directly or by railroading it over the bougie
10. If using a cuffed tube, inflate the cuff with air
11. Commence ventilation
12. Confirm correct placement of the tube by observation of movement of the chest, auscultation, and end-tidal CO₂ if available
13. Secure the tracheostomy tube by suturing it to skin and/or tracheal tape secured around the neck (Figure 17)



Figure 17: Tracheostomy tube secured with Velcro tape

Percutaneous Cricothyroidotomy using Seldinger Technique

Percutaneous cricothyrotomy using a Seldinger technique requires a needle with dilator and guide wire (Figure 18).




Figure 18: Cricothyroidotomy with Seldinger technique: guide wire (a); dilator (b); tracheostomy tube (c) (adapted from ⁴)

Percutaneous cricothyroidotomy: Surgical steps

1. Position the patient supine with neck exposed and extended (if possible)
2. Identify surface landmarks *i.e.* thyroid cartilage, cricoid cartilage and cricothyroid membrane
3. Prepare a sterile field
4. Inject 1% lidocaine with 1:100 000 epinephrine into the skin and through the cricothyroid membrane into the airway to anaesthetise the airway and suppress the cough reflex (if time to do so)
5. Fix the thyroid cartilage with the 1st & 3rd fingers of the non-dominant hand leaving the 2nd finger free to locate the cricothyroid membrane
6. With the dominant hand, make a small stab incision in the skin with a scalpel over the cricothyroid membrane (Figure 19)




Figure 19: Stab incision over cricothyroid membrane ⁴

7. Pass a finder needle attached to a syringe filled with normal saline, through the cricothyroid membrane, directing it caudally at 45⁰ (Figure 20).
8. Apply negative pressure to the syringe as the needle is advanced. Air bubbles will appear in the fluid-filled syringe as the needle traverses the membrane and enters the trachea.
9. Disconnect the syringe from the finder needle and insert the guide wire through the needle (Figure 21)




Figure 20: Pass needle through cricothyroid membrane⁴




Figure 21: Insert guide wire through the needle⁴

10. Retract and remove the needle once the guide wire has been advanced into the airway (Figure 22)




Figure 22: Remove the needle leaving guide wire in place⁴

11. Pass the dilator and tracheostomy tube over the guide wire
 12. Jointly advance the dilator and tracheostomy tube over the guide wire into the airway (Figure 23)




Figure 23: Jointly advance dilator and tracheostomy tube over guide wire⁴

13. Remove both dilator and guide wire leaving the tracheostomy tube *in situ* (Figure 24)




Figure 24: Tracheostomy tube in place⁴

14. Secure the tracheostomy tube with tracheostomy tape

Early Complications

- Bleeding
- Paratracheal false tract: Inadvertent extratracheal placement of the tracheostomy tube can be fatal. It is recognised by the absence of breath sounds on auscultation of the lungs, high ventilatory pressures, failure to ventilate the lungs, hypoxia, absence of expired CO₂, surgical emphysema, and an inability to pass a suction catheter down the bronchial tree, and on chest X-ray
- Posterior tracheal wall perforation into oesophagus
- Pneumothorax, surgical emphysema
- Hypercarbia and barotrauma

Late Complications

- Glottic or subglottic stenosis due to perichondritis and fibrosis of cricoid
- Dysphonia
- Persistent stoma
- Tracheoesophageal fistula

Postoperative care

Pulmonary oedema: This may occur following sudden relief of airway obstruction and reduction of high intraluminal airway pressures. It may be corrected by CPAP or positive pressure ventilation.

Respiratory arrest: This may occur immediately following insertion of the tracheostomy tube, and is attributed to the rapid reduction in arterial pCO₂ following restoration of normal ventilation, and hence loss of respiratory drive.

Humidification: A tracheostomy bypasses the nose and upper aerodigestive tract which normally warms, filters, and humidifies inspired air. To avoid tracheal desiccation and damage to the respiratory cilia and epithelium, and obstruction due to mucous crusting, the tracheostomy patient needs to breathe humidified warm air by means of a humidifier, heat and moisture exchange filter, or a tracheostomy bib.

Pulmonary Toilette: The presence of a tracheostomy tube and inspiration of dry air irritates the mucosa and increases secretions. Tracheostomy also promotes aspiration of saliva and food as tethering of the airway prevents elevation of the larynx during swallowing. Patients are unable to clear secretions as effectively as tracheostomy prevents generation of subglottic pressure, hence making coughing and clearing secretions ineffective; it also disturbs ciliary function. Therefore secre-

tions need to be suctioned in an aseptic and atraumatic manner.

Cleaning tube: Airway resistance is related to the 4th power of the radius with laminar flow, and the 5th power of the radius with turbulent flow. Therefore even a small reduction of airway diameter and/or conversion to turbulent airflow as a result of secretions in the tube can significantly affect airway resistance. Therefore regular cleaning of the inner cannula is required using a pipe cleaner or brush.

Securing tube: Accidental decannulation and failure to quickly reinsert the tube may be fatal. This is especially problematic during the 1st 48hrs when the tract has not matured and attempted reinsertion of the tube may be complicated by the tube entering a false tract. Therefore the tightness of the tracheostomy tapes should be regularly checked.

Cuff pressure: When tracheostomy tube cuff pressures against the tracheal wall mucosa exceed 30cm H₂O, mucosal capillary perfusion ceases, ischaemic damage ensues and tracheal stenosis may result. Mucosal injury has been shown to occur within 15 minutes. Therefore cuff inflation pressures of >25cm H₂O should be avoided. A number of studies have demonstrated the inadequacy of manual palpation of the pilot balloon as a means to estimate appropriate cuff pressures.

Measures to prevent cuff-related injury include:

- *Only inflate the cuff if required* (ventilated, aspiration)
- *Minimal Occluding Volume Technique:* Deflate the cuff, and then slowly reinflate until one can no longer hear air going past the cuff with a stethoscope applied to the side of the neck near the tracheostomy tube (ventilated patient)

- *Minimal Leak Technique:* The same procedure as above, except that once the airway is sealed, slowly to withdraw approximately 1ml of air so that a slight leak is heard at the end of inspiration
- *Pressure gauge:* Regular or continuous monitoring of cuff pressures

References

1. Dover K, Howdieshell TR, Colborn GL. The dimensions and vascular anatomy of the cricothyroid membrane: relevance to emergent surgical airway access. *Clin Anat.* 1996;9(5):291-5
2. Bennett JD, Guha SC, Sankar AB. Cricothyrotomy: the anatomical basis. *J R Coll Surg Edinb.* 1996 Feb;41(1): 57-60
3. Little CM, Parker MG, Tarnopolsky R. The incidence of vasculature at risk during cricothyroidostomy. *Ann Emerg Med.* 1986 Jul;15(7):805-7
4. Dept. Anaesthesia & Intensive Care, The Chinese University of Hong Kong: <http://www.aic.cuhk.edu.hk/web8/cricothyroidotomy.htm>

Author & Editor

Johan Fagan MBChB, FCORL, MMed
 Professor and Chairman
 Division of Otolaryngology
 University of Cape Town
 Cape Town, South Africa
johannes.fagan@uct.ac.za

THE OPEN ACCESS ATLAS OF OTOLARYNGOLOGY, HEAD & NECK OPERATIVE SURGERY

www.entdev.uct.ac.za



The Open Access Atlas of Otolaryngology, Head & Neck Operative Surgery by [Johan Fagan \(Editor\)](mailto:johannes.fagan@uct.ac.za) johannes.fagan@uct.ac.za is licensed under a [Creative Commons Attribution - Non-Commercial 3.0 Unported License](https://creativecommons.org/licenses/by-nc/3.0/)