OPEN ACCESS ATLAS OF OTOLARYNGOLOGY, HEAD & NECK OPERATIVE SURGERY

THE PECTORALIS MAJOR FLAP

The *pectoralis major flap* comprises the pectoralis major muscle, with or without overlying skin, and may include the underlying rib. It has an axial blood supply and is based superiorly on the pectoral branch of the thoracoacromial artery.

It is very useful in the head and neck, and can *inter alia* be used for the following: reconstruction of soft tissue defects of the oropharynx, oral cavity, hypopharynx, and skin of the neck; to augment pharyngeal repairs following salvage laryngectomy after previous chemoradiotherapy, and to cover carotid or jugular vein blow-outs *etc*. Rib may be included to bridge mandibular defects.

Relevant Anatomy

Pectoralis major muscle (Figure 1)

This muscle originates from the anterior aspect of the medial half of the clavicle; from the anterior surface of the sternum; from the cartilages of all the true ribs and from the aponeurosis of the abdominal external oblique muscle. The muscle fibers converge toward its insertion on the humerus. Muscle fibres arising from the clavicle pass transversely and are often separated from the rest of the muscle by a slight gap. The remainder of the muscle courses superolaterally.

Deep relations of pectoralis major muscle (Figure 2)

Deep to the pectoralis major muscle is its vascular pedicle, the pectoralis minor muscle, the costal cartilages, and inferiorly the costal attachments of the external oblique muscle.



Figure 1: Superficial dissection



Figure 2: Deep relations of pectoralis major muscle

Blood supply of pectoralis major flap (Figures 2, 3)

The pectoralis major flap is an axial flap and is based primarily on the pectoral branch of the thoracoacromial artery and its accompanying veins. The thoracoacromial artery is a branch of the axillary artery, itself a continuation of the subclavian artery. The pectoral branch of the thoraco-

Johan Fagan



Figure 3: Blood supply of pectoralis major flap

acromial artery courses within a well defined fascial plane on the deep surface of the pectoralis major muscle (Figures 2, 9). There is a clear dissection plane between this fascial layer and the superficial aspect of the pectoralis minor muscle making it possible to strip the pectoralis major and its blood supply off the pectoralis minor with blunt finger dissection. Additional blood supply arises medially from the internal mammary artery, and laterally from the long thoracic artery, branches of which are generally sacrificed during elevation of the flap to secure adequate pedicle length. Care must be taken should the skin island be located distally in the region of the costal margin at, or beyond, the lower extent of the pectoralis major muscle, as the blood supply then becomes random as opposed to axial, making perfusion of the flap less reliable.

Innervation

The pectoralis major is innervated by the lateral pectoral nerve which can be located just inferior to the clavicle with the pectoral branch of the thoracoacromial artery. The medial pectoral nerve passes through the pectoralis minor muscle and sends two to three branches to the pectoralis major. All these nerves are normally divided during elevation of the flap. The denervated muscle loses bulk with time which might be advantageous or disadvantageous, depending on the functional and cosmetic objectives of the flap.

Flap design

The flap may be employed either as a muscular or musculocutaneous flap, with or without the 4^{th} or 5^{th} ribs. The following discussion will focus on raising a musculocutaneous island flap.

Positioning, prepping and draping

The patient is placed in a supine position with the chest exposed and prepped up to the midline, and inferiorly to the costal margin. The upper arm is abducted slightly to expose the anterior axillary fold and lateral chest wall.

Surface markings of vascular pedicle

The surface markings of the vascular pedicle are determined by drawing a line from the shoulder to the xiphisternum and another line vertically from the midpoint of the clavicle to intersect the 1st line (*Figures* 4a, b).





Figures 4a,b: Surface markings of pectoral branch of thoracoacromial artery

Skin paddle design

The skin paddle is positioned over the pectoralis major muscle along the course of the pectoral branch of the thoracoacromial artery (*Figures 4, 5, 6*). In order to ensure that the pedicle is of adequate length, the distance between the top of the skin paddle and the inferior edge of the clavicle should equal or exceed the distance between the recipient site for the flap and the inferior edge of the clavicle. In women the paddle may be placed in the inframammary crease to include skin on either side of the crease to avoid excessive bulk from breast tissue and for cosmetic reasons (*Figure 5*).

Should additional pedicle length be required, the flap may be extended up to 2.5cms beyond the pectoralis major muscle inferiorly at the costal margin, with the knowledge that the flap then becomes a random pattern flap and the blood supply more tenuous.

Elevation of skin paddle

The skin is incised around the skin paddle, and the dissection is extended onto the sur-



Figure 5: Inframammary skin paddle

face of the pectoralis major muscle. As the vascular pedicle is located deep to the muscle, this may be quickly and safely performed using the bovie / monopolar diathermy. Care must be exercised not to undercut the skin paddle, but rather to bevel the dissection radially to include as many myocutaneous perforators as possible that supply the skin paddle (*Figure 6*). The skin paddle is tacked to the underlying pectoralis major muscle with a few sutures to minimise the risk of a shearing injury to the myocutaneous perforators (*Figure 7*).

Exposure of pectoralis major muscle

An incision is extended laterally from the peripheral margin of the skin paddle along the anterior axillary fold, which corresponds with the lateral margin of the pectoralis major muscle (*Figure 6*). The skin and breast tissue above the skin paddle are then widely elevated from the pectoralis major muscle with diathermy up to the clavicle (*Figure 7*).

Elevation of pedicle

The pectoralis major muscle is incised with cautery medially and inferiorly to the skin



Figure 6: Dissection of skin paddle onto pectoralis major muscle, and incision along anterior axillary fold



Figure 7: Exposure of pectoralis major muscle pedicle

paddle and is dissected from the ribs and intercostal muscles. The muscle is *not* divided superior to the skin paddle, as this might divide the vascular pedicle.

The pectoralis major muscle is then freed alongside the sternum with cautery. During this dissection perforators from the internal mammary artery are transected and cauterised.

The dissection plane between the pectoralis minor and major muscles and the vascular pedicle is found by dissecting along the lateral border of the pectoralis major muscle with electrocautery (*Figure 8*). Once this intermuscular plane has been identified, one can readily free the pectoralis major and its vascular pedicle from the pectoralis minor by stripping with a finger towards the clavicle (*Figure 8*).



Figure 8: Elevating pectoralis major from pectoralis minor along lateral margin of muscle

This brings the vascular pedicle (pectoral branch of thoracoacromial artery) clearly into view within the fascia on the deep surface of the pectoralis major muscle (*Figure 9*). The pedicle is kept in view and protected from injury while the branches from the lateral thoracic artery, and the branches of the medial pectoral nerve that traverse the pectoralis minor and enter the deep surface of the flap, are divided. The pectoralis major muscle is divided lateral to the pedicle while keeping the pedicle in view, thereby freeing it from the humerus.



Figure 9: Vascular pedicle visible on deep aspect of pectoralis major muscle

Skin tunnel over clavicle

The flap is generally passed into the neck superficial to the clavicle through a wide subcutaneous tunnel (*Figure 10*). The tunnel should be large enough to permit easy delivery of the flap into the neck without shearing the musculocutaneous perforators supplying the skin paddle, and to avoid strangulating the vascular pedicle. Dividing the subdermal connective tissue fibers of the skin above the skin tunnel with an upturned scalpel is a helpful adjunct to gain additional space.

After passing the flap into the neck, check that the pedicle is not twisted (*Figure 11*).

Gaining additional length

If the pedicle is too short, two manoeuvres may add additional length. The 1^{st} is to transect the pectoralis major muscle just below the clavicle, taking great care to preserve the vascular pedicle (*Figure 12*).



Figure 10: Flap is passed over the clavicle though a wide skin tunnel



Figure 11: Check that pedicle is not twisted



Figure 12: Vascular pedicle visible following division of muscular pedicle

An infraclavicular segment of muscle may also be excised. This has the benefit that the pedicle is directly applied to the clavicle, without intervening muscle. It has the added bonus of avoiding an unsightly, bulky pedicle over the clavicle.

A 2^{nd} manoeuvre to gain additional length is to pass the flap behind the clavicle (*Figure 13*).



Figure 13: Gaining additional length by passing pedicle behind clavicle

Should there be a concern about injury to the vascular structures deep to the clavicle, one can create a subperiosteal tunnel by incising the clavicular periosteum longitudenally along both the superior and inferior aspects of the clavicle. Since the periosteum is unyielding by its nature, this tunnel may be overly tight, but it can be relaxed by making incisions through the periosteum at its medial and distal ends, allowing easy passage of the flap. This manoeuvre adds some 3-4cm to the reach of the flap which may be just enough in a tight situation.

Closure of donor site defect

The donor site is either closed primarily with a closed suction drain, or a split skin graft is applied. Primary closure may be facilitated by undermining the surrounding skin.

Some Clinical examples



Figure 14: Reconstruction following total glossectomy



Figure 15: Augmentation of pharynx with pectoralis major myocutaneous flap following total laryngectomy with partial pharyngectomy



Figure 16: Reconstruction following segmental mandibulectomy and resection of tonsil, soft palate, and base of tongue



Figure 17: Laryngopharyngoesophagectomy defect reconstructed with tubed pectoralis major flap interposed between oropharynx and gastric pull-up that was too short to reach oropharynx





Figure 18a, b: Pectoralis major osteomyocutaneous flap used as salvage procedure following segmental mandibulectomy for osteoradionecrosis



Figure 19: Reconstruction of (a) pharyngocutaneous fistula with (b) pectoralis major muscle flap and (c) split skin graft and (d) final result

How to cite this chapter

Fagan JJ. (2025). Pectoralis major flap. In *The Open Access Atlas of Otolaryngology, Head & Neck Operative Surgery.* Retrieved from <u>https://vula.uct.ac.za/access/content/user/0</u> <u>1372298/Pectoralis%20major%20flap-</u> <u>1.pdf</u>

Flaps described in *The Open Access* Atlas of Otolaryngology Head & Neck Operative Surgery

- Pectoralis major flap
- <u>Cervicofacial flaps</u>
- <u>Deltopectoral flap</u>
- Buccal fat pad flap
- <u>Buccinator myomucosal flap</u>
- <u>Nasolabial flap</u>
- <u>Temporalis muscle flap</u>
- <u>Submental Artery Island flap</u>
- <u>Supraclavicular flap</u>
- Upper and lower trapezius flaps
- Latissimus dorsi flap
- Paramedian forehead flap
- Local flaps for facial reconstruction
- <u>Radial free forearm flap</u>
- <u>Anterolateral thigh (ALT) free flap</u>
- <u>Rectus abdominis flap</u>
- <u>Free fibula flap</u>
- <u>Thoracodorsal artery scapular tip</u> (TDAST) flap
- <u>Medial sural artery perforator (MSAP)</u> <u>flap</u>
- <u>Gracilis microvascular flap</u>
- <u>Principles and technique of</u> <u>microvascular anastomosis for free</u> <u>tissue transfer flaps in head and neck</u> <u>reconstructive surgery</u>

Author & Editor

Johan Fagan MBChB, FCS (ORL), MMed Emeritus Professor and Past Chair Division of Otolaryngology University of Cape Town Cape Town, South Africa johannes.fagan@uct.ac.za

THE OPEN ACCESS ATLAS OF OTOLARYNGOLOGY, HEAD & NECK OPERATIVE SURGERY www.entdev.uct.ac.za



The Open Access Atlas of Otolaryngology, Head & Neck Operative Surgery by <u>Johan Fagan (Editor)</u> johannes.fagan@uct.ac.za_is licensed under a <u>Creative</u> <u>Commons Attribution - Non-Commercial 3.0 Unported License</u>



