



The treatment of cancer of the supraglottic larynx is in a continual state of evolution. Clinicians are anxious to provide patients with the best possible functional outcome while controlling tumour. Therapeutic options for patients with cancer limited to the supraglottic region of the larynx include transoral microsurgical resection, chemoradiation therapy, open horizontal supraglottic laryngectomy and supracricoid laryngectomy.

Supraglottic laryngectomy was first introduced to the United States of America over 50 years ago. It effectively controls limited cancer involving the supraglottic portion of the larynx. The overwhelming majority of patients undergoing supra-glottic laryngectomy can swallow by mouth 10-14 days after surgery. However, the risk of aspiration pneumonia is significant in patients with inadequate cough and poor pulmonary reserve. It is therefore not appropriate for patients with cardiopulmonary compromise which is so significant that they cannot temporarily tolerate moderate amounts of aspiration. This group of patients should not be offered this procedure.

Patient Assessment

Tumour characteristics

All patients with cancer of the supraglottic larynx require careful pretreatment evaluation which should include laryngoscopy as well as imaging. The ideal candidates are those patients with tumour confined to the supraglottic larynx without involvement of either arytenoid. Cervical metastatic disease is highly prevalent for cancers of the supraglottic cancers. This risk is to both sides of the neck inasmuch as the larynx should be considered a midline structure. Routine bilateral functional neck dissection

of Levels 2, 3, and 4 is therefore advocated for all patients treated with surgery. Patients with bulky or bilateral metastases at presentation are considered a relative contraindication to supraglottic laryngectomy because of the predictable need for postoperative adjuvant radiation. Most patients can tolerate radiation therapy after supraglottic laryngectomy; however, it almost invariably compromises function with increased risk of aspiration.

Resection of tumour extension from the aryepiglottic fold to the medial wall of the pyriform sinus can be accomplished with modifications to become a partial laryngopharyngectomy. Tumour extension to the vallecula and tongue base can be accommodated, but this kind of tumour involvement will require partial tongue base resection, which in turn, will compromise functional outcome. Additionally, tongue base involvement almost always will call for postoperative irradiation, which, once again, is poorly tolerated after supraglottic laryngectomy.

The *only absolute contraindication* to supraglottic laryngectomy is spread through the paraglottic space inferiorly, such that the plane of resection through the ventricles would result in residual tumour or spread to involve both arytenoids. Under these circumstances, therapeutic alternatives include chemoradiation therapy or total laryngectomy.

Patient physiologic considerations

The patient's cardiopulmonary function is an essential criterion which requires evaluation when considering supraglottic laryngectomy. All patients require temporary tracheotomy which allows the treatment team to provide suctioning and to help treat aspiration. However, patients

with poor pulmonary reserve should not undergo supraglottic laryngectomy. There are no concrete, objective guidelines for eligibility such as arterial blood gases or spirometry. In general, patients with good functional capacity such that they can walk up a flight of stairs comfortably, can tolerate a supraglottic laryngectomy. Patients who cannot ambulate will, in general, not tolerate supraglottic laryngectomy.

Operative steps

The procedure is done under general anaesthesia. A prophylactic antibiotic is administered.

The patient is positioned in the supine position with the neck slightly extended on a shoulder roll. A superiorly based apron flap is planned such that the horizontal portion of the apron overlies the second or third tracheal ring (*Figure 1*). The **apron flap** is elevated just underneath the platysma muscle.




Figure 1: Supraglottic laryngectomy is accomplished through a superiorly based apron flap. The inferior aspect of the flap should not be extended past the midclavicular line because subsequent ischaemia may be associated with wound dehiscence

Tracheotomy is then accomplished. Under most circumstances this is facilitated by division and ligation of the thyroid isthmus.

Bilateral neck dissection is done in all patients. In the absence of identifiable pathologic adenopathy, elective neck dissection should remove the fatty lymphatic aponeurosis in Levels 2, 3 and 4. Level 2b superior to the spinal accessory nerve need not be dissected. The spinal accessory nerve, jugular veins, and sternocleidomastoid muscles are preserved.

The **suprahyoid strap muscles** are released from the superior surface of the hyoid bone with diathermy. The infrahyoid strap muscles are subsequently released approximately 1cm below the inferior insertion of the muscles to the hyoid bone and reflected inferiorly to expose the thyroid cartilage.

The **external perichondrium** of the thyroid cartilage is then released along the superior and lateral aspects of the thyroid laminae such that it can be reflected inferiorly (*Figure 2*).




Figure 2: The external perichondrium of the thyroid cartilage is incised along its superior insertion and reflected intact as an inferiorly based flap

The **cartilage** is then cut horizontally with an oscillating saw (*Figure 3*), recalling that

the intention is to resect the larynx through the ventricle (*Figure 4*). This requires that the cut be made approximately midway between the thyroid notch and the lower border of the thyroid cartilage. The cartilage cut may be slanted laterally toward the superior cornu on the less involved side which allows for preservation of the constrictor insertion.




Figure 3: The cartilage incision is carried horizontally midway between the thyroid notch superiorly and the inferior border of the thyroid cartilage in the midline to include the greater cornu of the thyroid cartilage on the side of the lesion. The greater cornu may be preserved on the side contralateral to the lesion or may be excised according to the judgment of the surgeon

Pharyngotomy is done through the vallecula when tumour is confined to the supraglottic larynx. This allows subsequent mucosal cuts to be made under direct vision.

Mucosal incisions are made laterally along the aryepiglottic folds and then across the face of the arytenoids to resect the entire epiglottis and false vocal folds. The inferior cuts are made horizontally through the laryngeal ventricles to connect the mucosal cuts with a cartilage cut (*Figure 4*).

The resected specimen is carefully evaluated and frozen sections done to assure complete tumour removal (*Figure 5*).

Haemostasis must be complete. This may be especially difficult along the cut edge of the tongue base where intermittent suture ligatures are recommended.




Figure 4: Mucosal cuts are made under direct vision. Inferiorly, the cut is made through the ventricle at the level of the true vocal cord. Posteriorly, the mucosal incision is made across the face of the arytenoid unless extension of tumour requires an extended procedure




Figure 5: Specimen includes epiglottis, hyoid bone, false vocal cords and upper half of thyroid cartilage

A **nasogastric tube** is inserted before the closure is accomplished.

Reconstruction starts with approximation of the tongue base to the glottic larynx in such a way that the tongue base is set back over the glottis to reduce the potential for aspiration (*Figure 6*). This requires that the yellow fibrofatty aponeurosis of the tongue base be directly approximated to the external perichondrium of the thyroid cartilage. If difficulty is observed with tearing of the external perichondrium, the sutures may be directed through residual thyroid cartilage. This initial layer of closure is then reinforced by approximating the suprahyoid musculature to the infrahyoid muscles. The closure is carried laterally to assure there is no evidence of salivary leak in the area of the pyriform sinus.





Figure 6: The external perichondrium of the thyroid cartilage is approximated to the fibroaponeurotic layer of the tongue base. This sets the tongue back over the glottis

After closure, **suction drains** are placed in either side of the neck and over the mucosal suture line and brought out through separate stab wounds.

The **apron flap is then redraped**. Great care is taken to assure that the **tracheotomy site is isolated** from the potential spaces in both sides following neck dissection by suturing (*Figures 7A & B*). Failure to isolate the tracheotomy from the potential space created by the neck dissection may result in transmission of contaminated mucus from the trachea underneath the

cervical flap and subsequent wound infection.

The anaesthesia tube is removed and replaced with a **cuffed tracheotomy tube**.



Figures 7A & B: At completion of the procedure, the tracheotomy site must be isolated from the potential dead space created through flap elevation and neck dissection

Postoperative Management

Care is taken to assure the integrity of the suction drains. Routine tracheotomy care is instituted. All patients are ambulated as soon as possible to assure restoration of tidal volume and prevention of atelectasis.

The **tracheotomy tube can be removed** when the patient demonstrates that he/she can protect the airway and is no longer aspirating. This usually takes 7-14 days. One practical method is to deflate the cuff of the tracheotomy tube when the patient is seated. Most patients will have some tem-

porary aspiration which can be suctioned. Thereafter it should be determined if the patient can tolerate an uninflated tracheotomy tube without excessive coughing and aspiration for 24hrs. Having achieved this milestone, the cuffed tracheotomy tube should be removed and replaced with a smaller tracheotomy tube without a cuff, which can be plugged so that the patient breathes past the tube and through the larynx, to ascertain if the patient's airway is satisfactory to allow decannulation.

The overwhelming majority of patients will tolerate an *oral diet* best when the tracheotomy site has healed. Introduction of an oral diet before the tracheotomy site is closed is difficult because the tracheotomy is associated with a less effective cough, tethering of the larynx, and diminished proprioception such that aspiration is more likely.

Following recovery from the acute phases of healing, supraglottic laryngectomy patients have remarkably good *voice* with few limitations to diet.

Recovery is significantly, and sometimes severely, challenged when *postoperative irradiation therapy* is required. Radiation causes lymphoedema and stasis, which compromises swallowing and may cause airway obstruction. Therapeutically induced obstructive sleep apnoea has been reported under these circumstances.

Patients who have had *prior irradiation* therapy can tolerate supraglottic laryngectomy. However, the healing is slower, recovery of deglutition is prolonged and some element of chondroradionecrosis of the residual thyroid cartilage may be observed. All these factors can be treated, and a successful functional outcome will be achieved in properly selected patients.

Cancer of Supraglottis: Clinical Practice Guidelines

<https://developingworldheadandneckcancerguidelines.com/index-page-supraglottic-cancers/>

Author

Jonas T. Johnson, M.D.
Dr Eugene N. Myers Professor & Chairman
Department of Otolaryngology
Eye and Ear Institute
200 Lothrop Street, Suite 500
Pittsburgh, PA 15213, USA
johnsonjt@upmc.edu

Editor

Johan Fagan MBChB, FCS(ORL), MMed
Professor and Chairman
Division of Otolaryngology
University of Cape Town
Cape Town, South Africa
johannes.fagan@uct.ac.za

THE OPEN ACCESS ATLAS OF OTOLARYNGOLOGY, HEAD & NECK OPERATIVE SURGERY

www.entdev.uct.ac.za



The Open Access Atlas of Otolaryngology, Head & Neck Operative Surgery by [Johan Fagan \(Editor\)](mailto:johannes.fagan@uct.ac.za) johannes.fagan@uct.ac.za is licensed under a [Creative Commons Attribution - Non-Commercial 3.0 Unported License](https://creativecommons.org/licenses/by-nc/3.0/)

