

## Air pollution and health

Bert Brunekreef, Stephen T Holgate

**The health effects of air pollution have been subject to intense study in recent years. Exposure to pollutants such as airborne particulate matter and ozone has been associated with increases in mortality and hospital admissions due to respiratory and cardiovascular disease. These effects have been found in short-term studies, which relate day-to-day variations in air pollution and health, and long-term studies, which have followed cohorts of exposed individuals over time. Effects have been seen at very low levels of exposure, and it is unclear whether a threshold concentration exists for particulate matter and ozone below which no effects on health are likely. In this review, we discuss the evidence for adverse effects on health of selected air pollutants.**

December, 2002, marks the 50th anniversary of the great smog event in London, UK. Stagnant weather conditions caused a sharp increase in the concentration of air pollutants, and over several days, more than three times as many people died than expected, leading to an estimated excess death toll of over 4000. Concentrations of sulphur dioxide and smoke reached several thousands of  $\mu\text{g per m}^3$ .<sup>1</sup> The London 1952 smog was not without precedent—similar events occurred in the Meuse valley, Belgium, in 1930, and elsewhere.<sup>2,3</sup>

Conditions have changed; effective legislation has eliminated most of the air pollution of 50 years ago. Yet the 1952 London smog event keeps attracting the attention of contemporary air pollution scientists. One question that remains important is the extent to which air pollution affects life expectancy. The 1954 report<sup>1</sup> suggests that death occurred mostly in individuals who were on the brink of death already, but if this were the case, the death rate should have dropped sharply after the episode. On the contrary, the death rate remained high for several months, and a recent re-analysis of the data indicates that the number of additional deaths due to the episode was about 12 000.<sup>4</sup> Another question is that of causality; although concentrations of sulphur dioxide and black smoke were greatly increased during the episode, sulphuric acid was thought to be the critical component. On this basis, bottles of ammonia were distributed among bronchitis patients so that they could neutralise acids during episodes of air pollution.<sup>5</sup>

In this review, we have focused on recent studies that have answered key questions that have arisen in the past 5 years. There is no shortage of recent reviews in this field,<sup>6–22</sup> and we have tried to add to, rather than duplicate, these reviews. For further reading, we have also added a list of relevant websites. These give access to comprehensive reviews produced by organisations such as the World Health Organisation, the European Union, the US Environmental Protection Agency, and the UK

Department for Environment, Food and Rural Affairs (panel).

### A new era of air pollution research

20 years ago, the era of successful abatement of traditional air pollutants culminated in a voluminous review of the health effects of ambient particulates.<sup>23</sup> At concentrations seen in the late 1970s in the developed world, adverse health effects were then regarded as unlikely. In the two decades since then, however, air pollution has re-emerged as a major environmental health issue. One reason is that, although air pollution from combustion of traditional fossil fuel is now present in much lower concentrations than 50 years ago, other components have gained prominence. Photochemical air pollution, characterised by high ozone concentrations during warm and sunny weather, was found to occur not only in places like Los Angeles and Mexico City, but also in large areas of Europe. Oxides of nitrogen produced by the ever rising number of motorised vehicles have increased until recently. Airborne particles have changed size distribution and composition, altering their toxicity.

#### *From episodes to time-series studies*

In the 1980s, a few air pollution episodes related to long-range transport from the east occurred in western Europe. In 1985, increases in mortality and hospital admissions in Germany and temporary lung function changes in the Netherlands were seen during one such episode.<sup>24,25</sup> Because concentrations of sulphur dioxide and particulate matter were in the hundreds rather than thousands of  $\mu\text{g/m}^3$  even under such episodic conditions, health effects of short episodes became hard to detect. Attention shifted to weather-driven, day-to-day variations in air pollution over long periods of time as determinants of day-to-day variations in mortality, hospital admissions, and other public-health indicators. Such time-series studies were

### Search strategy and selection criteria

The key words "air pollution and health" produce over 500 references a year from Medline alone. The present review is therefore, of necessity, our selection of just some of the major studies. The selection is based on a systematic Medline search until early 2002, on more than 20 years of continuous research in the field, and on participation in many advisory boards nationally and internationally.

*Lancet* 2002; **360**: 1233–42

**Institute for Risk Assessment Sciences, Utrecht University, PO Box 80176, 3508 TD Utrecht, Netherlands**  
(Prof B Brunekreef PhD); **and RCMB Division, School of Medicine, Southampton General Hospital, Southampton, UK**  
(Prof S T Holgate MD)

**Correspondence to:** Prof Bert Brunekreef  
(e-mail: B.Brunekreef@iras.uu.nl)

## Useful websites containing information on air pollution and health

### WHO

WHO air quality guidelines for Europe, 2000  
<http://www.euro.who.int/document/e71922.pdf>  
 WHO air quality guidelines for Europe, 2000 (background documents)  
<http://www.euro.who.int/air>  
 Transport, environment, and health, 2000  
<http://www.euro.who.int/document/e72015.pdf>

### US Environmental Protection Agency (EPA)

National ambient air quality standards (NAAQS), 2001  
<http://www.epa.gov/airs/criteria.html>  
 EPA's national ambient air quality standards: the standard review/re-evaluation process (1997)  
<http://www.epa.gov/ttn/oarpg/naaqsfm/naaqs.html>  
 Health and environmental effects of ground-level ozone (1997)  
<http://www.epa.gov/ttn/oarpg/naaqsfm/o3health.html>  
 Health and environmental effects of particulate matter (1997)  
<http://www.epa.gov/ttn/oarpg/naaqsfm/pmhealth.html>  
 Transportation and fuels (2002)  
<http://www.epa.gov/air/transport/index.html>  
 Air quality guide for ozone (1999)  
<http://www.epa.gov/airnow/consumer.html>  
 A guide to air quality and your health (2000)  
[http://www.epa.gov/airnow/aqi\\_cl.pdf](http://www.epa.gov/airnow/aqi_cl.pdf)  
 Air quality criteria for particulate matter (third external review draft), July 2002  
<http://cfpub.epa.gov/ncea/cfm/partmatt.cfm?ActType=default>

### European Union (EU)

European commission air quality website (2001)  
<http://europa.eu.int/comm/environment/air/>  
 Air quality framework directive (2002)  
<http://europa.eu.int/comm/environment/air/ambient.htm>  
 Clean air for Europe programme (2001)  
<http://europa.eu.int/comm/environment/air/cafe.htm>  
 EU Particulate Matter position paper (1997)  
[http://europa.eu.int/comm/environment/air/pp\\_pm.pdf](http://europa.eu.int/comm/environment/air/pp_pm.pdf)  
 EU ozone position paper (1999)  
[http://europa.eu.int/comm/environment/docum/pos\\_paper.pdf](http://europa.eu.int/comm/environment/docum/pos_paper.pdf)

### UK Department for Environment, Food and Rural Affairs

Air quality—what it means for your health (2001)  
<http://www.defra.gov.uk/environment/airquality/airpoll/index.htm>  
 Expert panel on air quality standards. Airborne particles: what is the appropriate measurement on which to base a standard? A discussion document (2001; 110 pp)  
[http://www.defra.gov.uk/environment/airquality/aqs/air\\_measures/index2.htm](http://www.defra.gov.uk/environment/airquality/aqs/air_measures/index2.htm)

### Other sources of information

A thematic network on air pollution and health: funded by EU, gives access to a network of research projects and information on air pollution and health.  
<http://airnet.iras.uu.nl>  
 The *Health and Clean Air Newsletter* is an attempt to make scientific information available to non-specialist readers, including reporters, without sacrificing accuracy.  
<http://healthandcleanair.org>

sporadic until about 1990,<sup>26,27</sup> but have increased exponentially since then. These studies have several advantages: weather-driven variations in air pollution concentrations generate large contrasts in exposure over time; populations serve as their own controls; and studies can often use routinely collected data. As a result, the number of deaths or hospital admissions studied can easily be in the hundreds of thousands, leading to great statistical power to detect small increases in adverse health effects of air pollution. Limitations of this approach include lack of individual characteristics; an assumption that exposure is representative for large populations by measurements made at central monitoring sites; and an assumption that confounding by long-term time trends, seasonal variations, weather, and influenza epidemics are controlled for by sophisticated statistical methods.

### Life shortening due to air pollution

Interest in health effects of air pollution became more intense after two US cohort studies suggested that exposure to fine particulate matter in the air was associated with life shortening.<sup>28,29</sup> Both studies were based on observations from the late 1970s to late 1980s, when air pollution concentrations were much lower than they had been in the past. A third cohort study (AHSMOG) found significant effects of particulate matter with a diameter of less than 10 µm (PM10) on non-malignant respiratory deaths in men and women, and on lung-cancer mortality in male, non-smoking Seventh-Day Adventists.<sup>30</sup> The effect on shortening life expectancy has been estimated at 1–2 years for realistic exposure contrasts,<sup>31</sup> which is substantial compared with the effects of other lifestyle or environmental risk factors related to mortality.

New observations from the largest study to date show that the findings persist after inclusion of several more

years of observation, with more consistent effects on lung cancer, in addition to non-malignant cardiopulmonary deaths, than in the original paper.<sup>32</sup> Case-control studies also continue to provide evidence of a link between air pollution (especially from traffic) and lung cancer.<sup>33</sup> Recent work has suggested that effects on life expectancy are not uniformly distributed but depend on factors such as education and antioxidant vitamin status,<sup>30,32,34</sup> which implies that life expectancy could be reduced more in disadvantaged population groups. On the basis of the cohort study findings, stringent standards for fine particulate matter have been proposed in the USA. Because large investments would be needed to improve air quality, the findings of the US cohort studies have received intense scrutiny (see section on disputing the evidence).

Until recently, no European cohort studies have provided data on life shortening. A Dutch study suggests that exposure to traffic-related air pollution is associated with cardiorespiratory deaths in much the same way as in the USA.<sup>35</sup>

### Pollutants of current interest: ozone, particulates, nitrogen dioxide

Now that the concentration of sulphur dioxide has decreased strikingly, attention has shifted to ozone, nitrogen dioxide, and particulates. Before discussing these pollutants in more detail, some qualification is needed to put our discussion in a wider global perspective. For millions of people living in rural areas in developing countries, indoor pollution from the use of biomass fuels occurs at concentrations that are orders of magnitude higher than currently seen in the developed world.<sup>36–38</sup> Deaths due to acute respiratory infections in children resulting from these exposures are estimated to be over

2 million per year. In the largest cities of the developing world, extreme exposures occur with both the traditional and the modern variety of pollutants.<sup>39,40</sup> Ideally, lessons learned in the developed world could help developing countries follow a more sustainable, less polluting path to industrialisation and modernisation. However, the available data suggest that the combined pressures of global competition and population increase leave little, if any, room to manoeuvre in this respect.

#### *Sources of ozone, particulate matter, and nitrogen dioxide*

Ozone is a strong oxidising agent formed in the troposphere through a complex series of reactions involving the action of sunlight on nitrogen dioxide and hydrocarbons.<sup>41</sup> Concentrations in city centres tend to be lower than those in suburbs, mainly as a result of the scavenging of ozone by nitric oxide originating from traffic.

The major source of anthropogenic emissions of nitrogen oxides into the atmosphere is the combustion of fossil fuels from stationary sources (heating, power generation) and in motor vehicles.<sup>41</sup> In ambient conditions, nitric oxide is rapidly transformed into nitrogen dioxide by atmospheric oxidants such as ozone.

Particulate air pollution is a mixture of solid, liquid, or solid and liquid particles suspended in the air. The size of suspended particles varies, from a few nm to tens of  $\mu\text{m}$ . The largest particles (coarse fraction) are mechanically produced by attrition of larger particles. Small particles ( $<1\ \mu\text{m}$ ) are largely formed from gases, the smallest ( $<0.1\ \mu\text{m}$ , ultrafine) of which are formed by nucleation resulting from condensation or chemical reactions that form new particles. In practical terms, a distinction is made between PM<sub>10</sub> ("thoracic" particles smaller than  $10\ \mu\text{m}$  in diameter that can penetrate into the lower respiratory system), PM<sub>2.5</sub> ("respirable" particles smaller than  $2.5\ \mu\text{m}$  that can penetrate into the gas-exchange region of the lung), and ultrafine particles smaller than  $100\ \text{nm}$  which contribute little to particle mass but which are most abundant in terms of numbers and offer a very large surface area, with increasing degrees of lung penetration.

### **Main findings from epidemiological studies**

#### *Short-term studies*

There have been abundant studies on the short-term effects of air pollution on health, with emphasis on mortality and hospital admissions. Panel studies have been done in volunteers, which have provided data on health endpoints such as respiratory and cardiovascular symptoms, and objective measures of lung or cardiac function on a daily or weekly basis. Large, collaborative efforts are under way in Europe and the USA which will be summarised.

In Europe, the APHEA (Air Pollution and Health: a European Approach) studies have provided many new insights. Initial studies were based on older data (APHEA-1),<sup>42</sup> and in the late 1990s a new series of studies (APHEA-2) was done which was able to make use of data on the PM<sub>10</sub> fraction. The APHEA-2 mortality study covered a population of more than 43 million people living in 29 European cities, which were all studied for more than 5 years in the early-mid 1990s.<sup>43</sup> From data involving 21 cities, the combined effect estimate showed that all-cause daily mortality increased by 0.6% (95% CI 0.4–0.8) for each  $10\ \mu\text{g}/\text{m}^3$  increase in PM<sub>10</sub>, with some heterogeneity to be discussed later. The APHEA-2 hospital admission study covered a population of 38 million living in eight European cities, which were studied for 3–9 years in the early to mid 1990s.<sup>44</sup> Hospital

admissions for asthma and chronic obstructive pulmonary disease (COPD) among people older than 65 years were increased by 1.0% (0.4–1.5) per  $10\ \mu\text{g}/\text{m}^3$  PM<sub>10</sub>,<sup>44</sup> and admissions for cardiovascular disease (CVD) were increased by about 0.5% (0.2–0.8) per  $10\ \mu\text{g}/\text{m}^3$  PM<sub>10</sub> and by about 1.1% (0.4–1.8) per  $10\ \mu\text{g}/\text{m}^3$  black smoke, suggesting an important role for diesel exhaust.<sup>45</sup> In APHEA-1, daily mortality increased in six cities<sup>46</sup> by 2.9% for each  $50\ \mu\text{g}/\text{m}^3$  increase in the 1-h maximum ozone concentration. Associations between nitrogen dioxide and mortality were also found, but these were sensitive to adjustment for black smoke, suggesting that the nitrogen dioxide represented a mixture of traffic-related air pollution. The corresponding data from APHEA-2 are awaited.

In the USA, the National Mortality, Morbidity and Air Pollution Studies (NMMAPS) focused on the 20 largest metropolitan areas in the USA, home to 50 million inhabitants, during 1987–94. All-cause mortality increased by 0.5% (0.1–0.9) for each  $10\ \mu\text{g}/\text{m}^3$  of PM<sub>10</sub>, a value close to the European results.<sup>47,48</sup> However, a recent communication suggests that an unrecognised software problem led to significant overestimation of this effect estimate, albeit without affecting its statistical significance or sensitivity to co-pollutant adjustment; details can be found at [http://www.healtheffects.org/Pubs/NMMAPS\\_letter.pdf](http://www.healtheffects.org/Pubs/NMMAPS_letter.pdf) (accessed July 11, 2002). Effects on hospital admissions were studied in ten cities with a combined population of 1 843 000 individuals older than 65 years.<sup>49</sup> Effects of PM<sub>10</sub> on COPD admissions were 1.5% (1.0–1.9) and on CVD admissions 1.1% (0.9–1.3) per  $10\ \mu\text{g}/\text{m}^3$  of PM<sub>10</sub>. There was also a weak effect of ozone on mortality in the summer but not winter. After adjustment for ozone and PM<sub>10</sub>, none of the other gaseous pollutants (including nitrogen dioxide) were associated with mortality.

Smaller-scale studies have been done on panels of individuals with daily or weekly observations of their exposure and health status.<sup>50,51</sup> These studies provide greater insight into the role of individuals' characteristics, but are dependent on participants' collaboration and provide relatively small data sets. These shortcomings make the results less reliable, especially with respect to uncontrolled confounding by incompletely modelled long-term time trends and medication use. One large-scale collaborative study, the PEACE (Pollution Effects on Asthmatic Children in Europe) study, has been completed in 28 regions of Europe. It failed to show effects of particulate matter and nitrogen dioxide on lung function and acute symptoms.<sup>52,53</sup> Because this was a winter-time study, no effects of ozone were assessed. Similar, smaller-scale studies have documented acute effects on lung function as well as acute symptoms in children and adults. One interpretation of these findings is that the inherent limitations of panel studies make them less suitable to detect subtle effects of air pollution than the large-scale mortality and hospital admission studies.

In both short-term and long-term studies, air pollution has an effect on cardiac deaths and hospital admissions in addition to respiratory effects. In Augsburg, Germany, during the 1985 air pollution episode, intriguing epidemiological observations were made serendipitously during a large cardiovascular study, the MONICA project. In retrospect, plasma viscosity, as well as heart rate and concentrations of C-reactive protein, were increased during the episode,<sup>54–56</sup> all of which can contribute to an increased risk of cardiovascular events. Studies in Boston, MA, USA, showed that nitrogen dioxide and PM<sub>2.5</sub> were associated with life-threatening arrhythmia leading to

therapeutic interventions by an implanted cardioverter defibrillator,<sup>37</sup> and that PM<sub>2.5</sub> concentrations were higher in the hours and days before onset of myocardial infarction in a large group of patients.<sup>38</sup> Cardiovascular events are now being studied in panel studies to ascertain more precisely the role of various pollutants and mechanisms.<sup>10</sup>

#### Long-term studies

In addition to cohort studies on mortality, air pollution effects on morbidity endpoints have been studied. Most of these have been cross-sectional, and assume that current air pollution exposure is sufficiently representative of long-term, previous exposure to make a plausible link with current health status. A series of Swiss studies on the association between air pollution and health among children and adults stand out for their careful design and conduct. Although Switzerland is a small country, exposure contrasts are relatively large because of the mountainous terrain. In eight different communities, lung function in adults was negatively associated with PM<sub>10</sub>, nitrogen dioxide, and sulphur dioxide,<sup>59</sup> in association with symptoms of bronchitis but not asthma.<sup>60</sup> In children from ten Swiss communities, the same pollutants were found to be associated with symptoms of bronchitis but not asthma or allergy.<sup>61</sup> The associations were seen at a range of PM<sub>10</sub> concentrations of 10–33 µg/m<sup>3</sup> only, which is well below the concentrations in many European countries. The high correlation between pollutants prevented separation of their individual effects.

In children living in 24 US and Canadian communities, significant associations were reported between exposure to fine particles and their acidity and lung function, symptoms of bronchitis, but again not asthma.<sup>62–64</sup> Exposure to particles has now also been related prospectively to reduced lung function growth in children,<sup>65</sup> and even children relocating from high to low pollution areas (or vice versa) were shown to experience changes in lung function growth that mirrored changes in exposure to particulate matter.<sup>66</sup>

Relation of variations in asthma prevalence to air pollution has been difficult;<sup>67,68</sup> however, prospective studies in California have recently suggested that some incident asthma cases<sup>69,70</sup> could be related to ozone but not other pollutants.

#### Mechanisms

Chamber studies provide a method by which to pursue the acute mechanisms of individual air pollutants, but do not reproduce either the mixtures or temporal variation that occur in natural exposures. Although individual air pollutants can exert their own specific individual toxic effects on the respiratory and cardiovascular systems, ozone, oxides of nitrogen, and suspended particulates all share a common property of being potent oxidants, either through direct effects on lipids and proteins or indirectly through the activation intracellular oxidant pathways.<sup>71</sup>

Animal and human in-vitro and in-vivo exposure studies have demonstrated the powerful oxidant capacity of inhaled ozone with activation of stress signalling pathways in epithelial cells<sup>72</sup> and resident alveolar inflammatory cells.<sup>73</sup> This mechanism involves activation of the transcription factor nuclear factor (NF) κB and its translocation to the nucleus. There it binds to DNA consensus sequences in the promoters of proinflammatory genes that code for cytokines (eg, granulocyte-macrophage colony-stimulating factor, tumour necrosis factor α [TNFα], and interleukin 1β), chemokines that attract neutrophils (eg, interleukin 8, neutrophil activating protein 78, and Gro α), and adhesion molecules (eg,

intercellular adhesion molecule 1).<sup>74</sup> These molecules increase neutrophil recruitment into the airways and alveoli and activate them for mediator secretion and the capacity to cause tissue damage.<sup>75,76</sup>

The large interindividual differences in responsiveness to inhaled ozone have an important genetic basis, as recently shown in mice<sup>77,78</sup> and human beings.<sup>79</sup> Candidate genes have included those for TNFα, manganese superoxide dismutase, glutathione peroxidase, NAD(P) quinone oxidoreductase, and glutathione S transferases. This finding emphasises the importance of locally available antioxidants such as uric acid, albumen, reduced glutathione, vitamin C, and vitamin E present in the lung lining fluid and epithelial barrier<sup>71,80</sup> in protecting the lung against ozone, and the protective effect exerted by diets supplemented with antioxidants.<sup>81</sup> An inflammatory response induced in the lung on exposure to ozone, together with increasing neuropeptide release from sensory neurons, contributes to the acute bronchoconstrictor response and hyper-responsiveness seen in asthma on exposure to this pollutant,<sup>82,83</sup> as well as to the tolerance induced by repeated short-term ozone exposure.<sup>83,84</sup> The significance of this induced tolerance on the recognised adverse effect of this pollutant on exacerbations of asthma during the summer months is not known. Studies have shown wide interindividual variability in responses to air pollutants. Although genetic factors undoubtedly account for part of this inconsistency, other more subtle factors could be operating, such as the distribution of ventilation in the different lung compartments, and in the case of particles, differences in regional deposition within the lungs.

By contrast with ozone, little is known about the effects of nitrogen dioxide on normal and diseased lung. In-vitro studies in animals and human beings confirm the capacity of nitrogen dioxide to activate oxidant pathways, albeit less potently than ozone.<sup>72,85</sup> The ensuing inflammatory response also differs by enhancing the recruitment of T lymphocytes and macrophages.<sup>86</sup> However, one feature of nitrogen dioxide that might contribute to exacerbations of respiratory disease is its capacity to impair the function of alveolar macrophages and epithelial cells, thereby increasing the risk of lung infection.<sup>87,88</sup> Although nitrogen dioxide can also enhance airway responses to inhaled allergens in asthmatic individuals,<sup>89</sup> short-term exposure remains relatively benign. Little is known about the long-term effects of nitrogen dioxide in human beings, but in rodents, prolonged exposure to either ozone or nitrogen dioxide results in destruction of peripheral airways.

The increase in respiratory and cardiovascular morbidity and mortality that has been epidemiologically linked to inhaled particulates has, until recently, defied a mechanistic explanation. In-vitro and in-vivo studies in animals and human beings have revealed potent proinflammatory effects involving lung epithelial cells<sup>90</sup> and alveolar macrophages.<sup>91</sup> Both directly and through uptake into epithelial cells<sup>92</sup> and macrophages,<sup>93,94</sup> oxidant pathways are activated<sup>95,96</sup> with the downstream consequences of stimulating cytokine and mediator release,<sup>97</sup> resulting in extensive neutrophil migration, but also T lymphocyte recruitment and activation.<sup>98,99</sup> On reaching the bone marrow, cytokines and chemokines released from the lung stimulate egression of neutrophils and their precursors into the circulation.<sup>100</sup> In the short term there is acute tissue damage with activation of the epidermal-growth-factor receptor (EGFR) pathway, and evidence for organ-repair responses.<sup>101</sup> This reaction is partly due to surface processing of EGFR ligands such as heparin-binding EGF-like growth factor (HB-EGF) and

oxidant-induced transactivation of the EGFR.<sup>102</sup> If this cycle of damage and repair continues, epithelial mucus metaplasia results, as does ongoing cytokine and chemokine secretion that contributes to airway inflammation.

Although knowledge of the respiratory actions of particulates has grown, an understanding of how particulates increase the risk of cardiovascular events has proven much more difficult to achieve. Again, through activation of stress signalling pathways from the epithelium to the lung microvessels, factors that influence blood clotting are generated. Increased concentrations of fibrinogen and platelets, and sequestration of red blood cells in the lung mass<sup>103</sup> have also been detected in relation to particulate pollution, but, along with increasing the risk of cardiac arrhythmia,<sup>10</sup> their significance in the cardiovascular events linked to particle pollution remains to be established.<sup>11</sup>

Diesel particulates and ozone have been shown to increase the synthesis of the allergic antibody IgE in animals<sup>104</sup> and human beings,<sup>105</sup> which would increase sensitisation to common allergens.<sup>106</sup> By interacting together and with other environmental factors, particulates and gaseous air pollutants can have long-term effects on allergic individuals. Although pollutants are unlikely to be able to interact to enhance proallergic and inflammatory responses, no studies have investigated mixtures. Similarly, long-term exposure to pollutant mixtures can have tissue-damaging effects which could be irreversible.<sup>13</sup> The recent recognition that ultrafine particles (mass median diameter <0.1 µm) are more toxic when inhaled than PM10 suggests that their ability to be absorbed into tissues and the circulation, and their greatly increased surface area, might be important factors in determining cardiopulmonary toxicity.<sup>107</sup>

### Air quality guidelines and standards

Several guidelines and standards exist for ozone, nitrogen dioxide, and particulate matter in ambient air. The table lists the most recent air quality guidelines and standards recommended by WHO, the US Environmental Protection Agency, and the European Union (EU). The EU standards are targets to be reached in 2005 or 2010. The most remarkable difference lies in the annual value for nitrogen dioxide. The WHO and EU value is only 40% of the US value.

WHO has not proposed guidelines for particulate matter, arguing that it was unable to define a threshold below which no adverse effects are expected. Instead, dose-response information was provided to help policy makers decide when setting a standard. The effect estimates given in the table were based on information as it was available until 1996. As described earlier, the newest large-scale studies<sup>43,48</sup> tend to show somewhat smaller effects per unit particulate matter. For particulate matter, the proposed US standards for PM<sub>2.5</sub> are not very different from the EU 2010 annual average and the 2005 24-h average for PM<sub>10</sub>, considering that PM<sub>2.5</sub> usually comprises about 60–70% of the PM<sub>10</sub> concentration, and considering the number of exceedances allowed in the 24-h EU standard.

All guidelines and standards mentioned in the table are subject to periodic revision when new scientific information becomes available. WHO has just recently started a process to re-evaluate the guidelines for these three pollutants.

### Current issues

#### Thresholds

A key question is whether threshold concentrations exist below which air pollution has no effect on population health. If such a threshold could be identified, no additional

	Maximum concentration allowed when averaged over time*			
	1 h	8 h	24 h	1 year
<b>Ozone (µg/m<sup>3</sup>)</b>				
WHO	..	120	..	..
EPA	235	157 (proposed)	..	..
EU	..	120 (2010)	..	..
<b>Nitrogen dioxide (µg/m<sup>3</sup>)</b>				
WHO	200	..	..	40
EPA	..	..	..	100
EU	200 (2010)	..	..	40 (2010)
<b>PM<sub>10</sub> (µg/m<sup>3</sup>)</b>				
WHO (mortality relative risk per 10 µg/m <sup>3</sup> )†	..	..	1.007	1.10
EPA	..	..	150	50
EU	..	..	50‡ (2005), 50§ (2010)	40 (2005), 20 (2010)
<b>PM<sub>2.5</sub> (µg/m<sup>3</sup>)</b>				
WHO (mortality relative risk per 10 µg/m <sup>3</sup> )†	..	..	1.015	1.14
EPA	..	..	65 (proposed)	15 (proposed)

For details see <http://www.euro.who.int/document/e71922.pdf>, <http://www.epa.gov/airs/criteria.html>, and <http://www.ircline.be>. \*Short averaging times are used when the guideline was developed to prevent acute effects, long averaging times to prevent long-term effects. †No guideline value for particulate matter was given because no threshold concentration was identified below which no effects on health were expected. Relative risk estimates were provided to help policy makers set standards based on quantitative dose-response information. ‡To be exceeded on no more than 35 days per year. §To be exceeded on no more than 7 days per year.

### WHO, US Environmental Protection Agency (EPA), and European Union (EU) air quality guidelines and standards for ozone, nitrogen dioxide, and particulate matter

public-health benefits would be expected from bringing air pollution concentrations far below this level. Theoretical and empirical work has been done to shed light on this issue.<sup>108,109</sup> In an analysis of NMMAPS data, no evidence was found for a threshold for PM<sub>10</sub> and daily all-cause and cardiorespiratory mortality.<sup>109</sup> By contrast, a threshold of about 50 µg/m<sup>3</sup> was estimated for non-cardiorespiratory causes of death, illustrating the specificity of the approach. Earlier analyses restricted the analysis to concentrations below a certain value.<sup>37,110</sup> These analyses suggest that a threshold for acute effects of ozone on lung function changes must lie well below 100 µg/m<sup>3</sup> as an hourly maximum.

#### Displacement of daily mortality and hospital admissions

Although time-series studies have shown a link between day-to-day variations in air pollution concentrations and daily deaths and hospital admissions, by how many days, weeks, or months such events are brought forward is unclear. If deaths occur just a few days earlier than they would have occurred anyway, the public-health significance of these associations would be much less than if life expectancy is being reduced by months or years.<sup>111</sup> Recent analyses show that, in the time-series studies, there is no evidence that deaths or hospital admissions are being brought forward by just a few days.<sup>112–115</sup> On the contrary, effect estimates increase with increasing duration of exposure to air pollution, suggesting that cumulative exposures have stronger effects on mortality than can be extracted from associations between day-to-day variations in air pollution and deaths. Previous data have shown that many of the deaths associated with air pollution were occurring outside hospital, which also supports the suggestion that these patients were often not terminally ill.<sup>116</sup>

### *Relation between ambient and personal exposure to particulate matter, nitrogen dioxide, and ozone in short-term and long-term studies*

In developed countries, people spend more than 80% of their time indoors, and most of that time in their own home. Indoor pollutant concentrations can differ from outdoor concentrations because of ventilation patterns and indoor sources. Therefore whether measurement of air pollution outdoors is a valid method of assessing exposure of the population has been questioned. The day-to-day variation in personal exposure to particulate matter correlates with the day-to-day variation in ambient concentrations of particulate matter; correlations are especially high in the case of fine particles (PM<sub>2.5</sub>).<sup>117</sup> One study that addressed this issue simultaneously for particulate matter and gaseous components found that there was no association between the day-to-day variation in personal exposure to nitrogen dioxide, sulphur dioxide, and ozone and ambient measurements of these three components.<sup>118</sup> Ambient PM<sub>2.5</sub>, nitrogen dioxide, sulphur dioxide, and ozone were closely associated with personal PM<sub>2.5</sub>, which strongly suggests that gaseous and PM<sub>2.5</sub> concentrations outdoors act as a surrogate for personal exposure to PM<sub>2.5</sub>.<sup>118</sup>

Few studies have addressed the question of whether spatial variations in long-term average outdoor concentrations are reflected in similar variations in long-term personal exposures. For PM<sub>2.5</sub> and nitrogen dioxide, regional and local-scale spatial variations in outdoor concentrations have been shown to be reflected in similar variations in personal exposures.<sup>119,120</sup> The US AHSMOG study and the recent Dutch cohort study have used estimates of small-scale spatial variations in air pollution exposure to their advantage,<sup>30,121</sup> suggesting that further improvements can be expected when such within-community differences in exposure are taken into account.

### *Causality*

Knowledge of which pollution components are responsible for any health effects observed in epidemiological studies is of obvious importance. For ozone, the situation is relatively simple. A large experimental database exists to document that ozone has significant biological effects at ambient concentrations. In addition, the correlation between ozone and other pollutant concentrations in outdoor air is often low, so the effects of ozone and other pollutants can be separated relatively easily.

For nitrogen dioxide, the situation is more complex. Evidence for biological effects at ambient concentrations is much weaker than for ozone. In outdoor air, nitrogen dioxide is often highly correlated with other combustion products, notably fine particulate matter, and to personal PM<sub>2.5</sub>, but not personal nitrogen dioxide.<sup>118</sup> Our assessment is that, in most circumstances, nitrogen dioxide serves as a surrogate for all traffic-related combustion products.

Most complex is the question of which particulate matter components or attributes are most important in determining health effects. Many candidates have been proposed. One is ultrafine particles—ie, particles of less than 100 nm which can be found in very high numbers (>100 000 cm<sup>-3</sup>) near busy roads.<sup>122</sup> The thinking is that such particles can easily find their way from the lungs into the bloodstream and then lead to systemic inflammatory changes which may affect blood coagulability. Similarly, some suggestive evidence from epidemiological studies points to ultrafine particles being related to respiratory

health endpoints.<sup>123</sup> However, the number of ultrafine particles in the air is often poorly correlated with PM<sub>2.5</sub>, so ultrafine particles are unlikely to explain much of the association between particulate matter mass and health endpoints.<sup>124</sup>

Specific components related to traffic exhaust, especially diesel combustion products, could be important.<sup>125</sup> The APHEA and NMMAPS analyses have suggested that particulate matter effects are larger in areas with high nitrogen dioxide (ie, traffic density)<sup>43</sup> or in areas with high emissions of particulate matter from highway vehicles and diesel locomotives.<sup>126</sup> An intriguing observation from a time-series study is that individuals who live on the main roads of Amsterdam have much higher relative risks of death than people who live away from the main roads, when analysed with data from the same background air pollution monitoring station.<sup>127</sup> This finding clearly suggests that traffic-related particulate matter is involved in explaining increases in mortality on high pollution days.

Factor analysis suggested that particles from mobile and coal combustion sources, but not coarse particles originating from the earth's crust, are responsible for effects on mortality.<sup>128</sup> Studies from the Utah Valley, USA, have linked transition metal content of particulate matter with particulate matter toxicity in human bronchial instillation studies.<sup>129</sup>

Much attention has been devoted to separating the effects of fine particulate matter (PM<sub>2.5</sub>) and coarse mass. Initial observations that fine particulate matter was associated with mortality whereas coarse mass was not<sup>130</sup> have been corroborated in several recent studies,<sup>131,132</sup> but others have found independent effects of coarse mass on hospital admissions for asthma,<sup>133</sup> and one study was unable to separate effects of fine particulate matter from those of coarse mass.<sup>134</sup>

Support for causality also comes from studies on effects of reducing air pollution on health outcomes. One of the best examples is a labour dispute that shut down a large steel mill in the Utah Valley for 14 months in 1987. Ambient particulate matter concentrations as well as respiratory hospital admissions were clearly decreased during the strike, only to increase to prestrike levels after the dispute ended.<sup>135</sup> Mortality was decreased as well.<sup>136</sup> In combination with the recent toxicological studies of particulate matter collected before, during, and after the strike, the Utah Valley example provides strong evidence of a causal relation between exposure to ambient particulate matter and mortality and morbidity. Other examples include reductions in acute-care visits and hospital admissions for asthma in Atlanta, GA, USA, in conjunction with reduced air pollution due to traffic measures taken during the 2000 Olympic games,<sup>137</sup> and reductions in bronchitis in association with reduced air pollution over several years in the former German Democratic Republic.<sup>138</sup>

### **Disputing the evidence**

In view of the potentially large costs and benefits associated with abatement of air pollution, questions surrounding the relation between air pollution and health have been an area of fierce debate in the past decade. The early time-series studies have been criticised for their analytical approach and inadequate control for confounding by weather variables etc, whereas the US cohort studies have been criticised for inadequate confounder and co-pollutant control. Reanalyses of such studies were done to investigate whether findings depended more on analytical approach than on robust

relations. The USA-based Health Effects Institute, itself a partnership between (automobile) industry and government, has taken a leadership role in this debate by commissioning two reanalysis projects—one to reanalyse the Philadelphia time-series study,<sup>139</sup> which had generated much of the debate on short-term effects, and another to reanalyse the two main US cohort studies. These reanalyses were done by independent researchers who were granted access to all original data and largely confirmed the findings of the original studies.<sup>140,141</sup> In addition, the reanalysis revealed new insights into the role of weather variables in time-series studies,<sup>142,143</sup> and new methods for enabling analysis of spatial association between air pollution, mortality, and potential confounding variables.<sup>144</sup> The reanalyses themselves have served as role models on how to provide policy makers with the best possible scientific evidence.<sup>145</sup>

### Concluding remarks

An excess risk of death of “0.5% per 10 µg/m<sup>3</sup> PM10” requires some translation before the effect on public health becomes clear. For the Netherlands (16 million inhabitants, about 140 000 deaths per year, and an average PM10 concentration of >30 µg/m<sup>3</sup>), the number of deaths attributable to day-to-day variations in PM10 would translate into at least 2100 deaths brought forward by air pollution per year—almost twice the number of deaths due to traffic accidents. Estimates derived from the cohort studies are much higher still because these incorporate long-term as well as short-term effects. For Austria, France, and Switzerland combined (population about 74.5 million), 40 000 deaths per year are estimated to be attributable to air pollution, about half to air pollution from traffic specifically.<sup>146</sup> Similarly high numbers have been estimated for respiratory and cardiovascular hospital admissions, bronchitis episodes, and restricted activity days. Because of such numbers, health effects from air pollution have been estimated to be higher than effects from a long list of other environmental factors.<sup>147</sup> These estimates are based on three major assumptions: causality of the epidemiological associations, linearity of the exposure-response relations, and that a threshold is absent or has a very low value. All of these assumptions are being tested rigorously; further proof could arise from research specifically targeted at assessment of changes in air pollution concentrations, such as those being done in a series of studies in the former German Democratic Republic.<sup>138</sup>

Given the high cost of further measures to reduce air pollution, and the many new findings which suggest that health effects can be seen at ever lower concentrations, the health effects of air pollution will need to receive much scientific and regulatory interest for years to come.

*Conflict of interest statement*  
None declared.

### References

- Ministry of Health. Mortality and morbidity during the London fog of December 1952. Reports on Public Health and Medical Subjects No 95. London: HMSO, 1954.
- Nemery B, Hoet PH, Nemmar A. The Meuse Valley fog of 1930: an air pollution disaster. *Lancet* 2001; **357**: 704–08.
- Firket J. Fog along the Meuse Valley. *Trans Faraday Soc* 1936; **32**: 1192–97.
- Bell ML, Davis DL. Reassessment of the lethal London fog of 1952: novel indicators of acute and chronic consequences of acute exposure to air pollution. *Environ Health Perspect* 2001; **109** (suppl 3): 389–94.
- Anon. An anti-smog bottle. *BMJ* 1955; **231**: 1135.
- Goldsmith CA, Kobzik L. Particulate air pollution and asthma: a review of epidemiological and biological studies. *Rev Environ Health* 1999; **14**: 121–34.
- Koenig JQ. Air pollution and asthma. *J Allergy Clin Immunol* 1999; **104**: 717–22.
- Nicolai T. Air pollution and respiratory disease in children: what is the clinically relevant impact? *Pediatr Pulmonol Suppl* 1999; **18**: 9–13.
- Pope CA III. Mortality and air pollution: associations persist with continued advances in research methodology. *Environ Health Perspect* 1999; **107**: 613–14.
- Donaldson K, Gilmour MI, MacNee W. Asthma and PM10. *Respir Res* 2000; **1**: 12–15.
- Gordon T, Reibman J. Cardiovascular toxicity of inhaled ambient particulate matter. *Toxicol Sci* 2000; **56**: 2–4.
- Nyberg BF, Pershagen G. Epidemiologic studies on the health effects of ambient particulate air pollution. *Scand J Work Environ Health* 2000; **26** (suppl 1): 49–89.
- Pope CA 3rd. Epidemiol fine particulate air Pollut Hum health: biologic mechanisms and who's at risk? *Environ Health Perspect* 2000; **108** (suppl 4): 713–23.
- Takafuji S, Nakagawa T. Air pollution and allergy. *J Invest Allergol Clin Immunol* 2000; **10**: 5–10.
- Dockery DW. Epidemiologic evidence of cardiovascular effects of particulate air pollution. *Environ Health Perspect* 2001; **109** (suppl 4): 483–86.
- Donaldson K, Stone V, Seaton A, MacNee W. Ambient particle inhalation and the cardiovascular system: potential mechanisms. *Environ Health Perspect* 2001; **109** (suppl 4): 523–27.
- Gilmour MI, Daniels M, McCrillis RC, Winsett D, Selgrade MK. Air pollutant-enhanced respiratory disease in experimental animals. *Environ Health Perspect* 2001; **109** (suppl 4): 619–22.
- Oberdorster G. Pulmonary effects of inhaled ultrafine particles. *Int Arch Occup Environ Health* 2001; **74**: 1–8.
- Peden DB. Air pollution in asthma: effect of pollutants on airway inflammation. *Ann Allergy Asthma Immunol* 2001; **87**: 12–17.
- Sunyer J. Urban air pollution and chronic obstructive pulmonary disease: a review. *Eur Respir J* 2001; **17**: 1024–33.
- Sydbom A, Blomberg A, Parnia S, et al. Health effects of diesel exhaust emissions. *Eur Respir J* 2001; **17**: 733–46.
- Thurston GD, Ito K. Epidemiological studies of acute ozone exposures and mortality. *J Expo Anal Environ Epidemiol* 2001; **11**: 286–94.
- Holland WW, Bennett AE, Cameron IR, et al. Health effects of particulate pollution: reappraising the evidence. *Am J Epidemiol* 1979; **110**: 527–659.
- Wichmann HE, Mueller W, Allhoff P, et al. Health effects during a smog episode in West Germany in 1985. *Environ Health Perspect* 1989; **79**: 89–99.
- Dassen W, Brunekreef B, Hoek G, et al. Decline in children's pulmonary function during an air pollution episode. *J Air Pollut Control Assoc* 1986; **36**: 1223–27.
- Ostro B. A search for a threshold in the relationship of air pollution to mortality: a reanalysis of data on London winters. *Environ Health Perspect* 1984; **58**: 397–99.
- Schwartz J, Marcus A. Mortality and air pollution in London: a time series analysis. *Am J Epidemiol* 1990; **131**: 185–94.
- Dockery DW, Pope CA 3rd, Xu X, et al. An association between air pollution and mortality in six US cities. *N Engl J Med* 1993; **329**: 1753–59.
- Pope CA 3rd, Thun MJ, Namboodiri MM, et al. Particulate air pollution as a predictor of mortality in a prospective study of US adults. *Am J Respir Crit Care Med* 1995; **151**: 669–74.
- Abbey DE, Nishino N, McDonnell WF, et al. Long-term inhalable particles and other air pollutants related to mortality in nonsmokers. *Am J Respir Crit Care Med* 1999; **159**: 373–82.
- Brunekreef B. Air pollution and life expectancy: is there a relation? *Occup Environ Med* 1997; **54**: 781–84.
- Pope CA 3rd, Burnett RT, Thun MJ, et al. Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *JAMA* 2002; **287**: 1132–41.
- Nyberg F, Gustavsson P, Jarup L, et al. Urban air pollution and lung cancer in Stockholm. *Epidemiology* 2000; **11**: 487–95.
- Brunekreef B. All but quiet on the particulate front. *Am J Respir Crit Care Med* 1999; **159**: 354–56.
- Hoek G, Brunekreef B, Goldbohm S, Fischer P, Brandt P. Association between mortality and indicators of traffic-related air pollution in the Netherlands: a cohort study. *Lancet* (in press).
- McMichael AJ, Smith KR. Seeking a global perspective on air pollution and health. *Epidemiology* 1999; **10**: 1–4.
- Smith KR, Samet JM, Romieu I, Bruce N. Indoor air pollution in developing countries and acute lower respiratory infections in children. *Thorax* 2000; **55**: 518–32.

- 38 Ezzati M, Kammen D. Indoor air pollution from biomass combustion and acute respiratory infections in Kenya: an exposure-response study. *Lancet* 2001; **358**: 619–24.
- 39 Vichit-Vadakan N, Ostro BD, Chestnut LG, et al. Air pollution and respiratory symptoms: results from three panel studies in Bangkok, Thailand. *Environ Health Perspect* 2001; **109** (suppl 3): 381–87.
- 40 Ostro B, Chestnut L, Vichit-Vadakan N, Laixuthai A. The impact of particulate matter on daily mortality in Bangkok, Thailand. *J Air Waste Manag Assoc* 1999; **49**: 100–07.
- 41 WHO. Air Quality Guidelines for Europe, 2nd edn. *WHO Reg Publ Eur Ser* 2000; **91**: 1–287.
- 42 Katsouyanni K, Zmirou D, Spix C, et al. Short-term effects of air pollution on health: a European approach using epidemiological time-series data. The APHEA project: background, objectives, design. *Eur Respir J* 1995; **8**: 1030–38.
- 43 Katsouyanni K, Touloumi G, Samoli E, et al. Confounding and effect modification in the short-term effects of ambient particles on total mortality: results from 29 European cities within the APHEA2 project. *Epidemiology* 2001; **12**: 521–31.
- 44 Atkinson RW, Anderson HR, Sunyer J, et al. Acute effects of particulate air pollution on respiratory admissions: results from APHEA 2 project. *Air Pollution and Health: a European Approach*. *Am J Respir Crit Care Med* 2001; **164**: 1860–66.
- 45 Le Tertre A, Medina S, Samoli E, et al. Short term effects of particulate air pollution on cardiovascular diseases in eight European cities. *J Epidemiol Community Health* (in press).
- 46 Touloumi G, Katsouyanni K, Zmirou D, et al. Short-term effects of ambient oxidant exposure on mortality: a combined analysis within the APHEA project. *Air Pollution and Health: a European Approach*. *Am J Epidemiol* 1997; **146**: 177–85.
- 47 Samet JM, Dominici F, Zeger SL, Schwartz J, Dockery DW. The National Morbidity, Mortality, and Air Pollution Study. Part I: methods and methodologic issues. *Res Rep Health Eff Inst* 2000; **75**: 5–14.
- 48 Samet JM, Dominici F, Currier FC, Coursac I, Zeger SL. Fine particulate air pollution and mortality in 20 US cities, 1987–1994. *N Engl J Med* 2000; **343**: 1742–49.
- 49 Zanobetti A, Schwartz J, Dockery DW. Airborne particles are a risk factor for hospital admissions for heart and lung disease. *Environ Health Perspect* 2000; **108**: 1071–77.
- 50 Pope CA 3rd, Dockery DW, Spengler JD, Raizenne ME. Respiratory health and PM10 pollution: a daily time series analysis. *Am Rev Respir Dis* 1991; **144**: 668–74.
- 51 Pope CA. 3rd, Dockery DW. Acute health effects of PM10 pollution on symptomatic and asymptomatic children. *Am Rev Respir Dis* 1992; **145**: 1123–28.
- 52 Roemer W, Hoek G, Brunekreef B, et al. Daily variations in air pollution and respiratory health in a multicentre study: the PEACE project. *Pollution Effects on Asthmatic Children in Europe*. *Eur Respir J* 1998; **12**: 1354–61.
- 53 Roemer W, Hoek G, Brunekreef B. Pollution effects on asthmatic children in Europe, the PEACE study. *Clin Exp Allergy* 2000; **30**: 1067–75.
- 54 Peters A, Doring A, Wichmann HE, Koenig W. Increased plasma viscosity during an air pollution episode: a link to mortality? *Lancet* 1997; **349**: 1582–87.
- 55 Peters A, Frohlich M, Doring A, et al. Particulate air pollution is associated with an acute phase response in men; results from the MONICA-Augsburg Study. *Eur Heart J* 2001; **22**: 1198–204.
- 56 Peters A, Perz S, Doring A, et al. Increases in heart rate during an air pollution episode. *Am J Epidemiol* 1999; **150**: 1094–98.
- 57 Peters A, Liu E, Verrier RL, et al. Air pollution and incidence of cardiac arrhythmia. *Epidemiology* 2000; **11**: 11–17.
- 58 Peters A, Dockery DW, Muller JE, Mittleman MA. Increased particulate air pollution and the triggering of myocardial infarction. *Circulation* 2001; **103**: 2810–15.
- 59 Ackermann-Lieblich U, Leuenberger P, Schwartz J, et al. Lung function and long term exposure to air pollutants in Switzerland. Study on Air Pollution and Lung Diseases in Adults (SAPALDIA) Team. *Am J Respir Crit Care Med* 1997; **155**: 122–29.
- 60 Zemp E, Elsasser S, Schindler C, et al. Long-term ambient air pollution and respiratory symptoms in adults (SAPALDIA study). *Am J Respir Crit Care Med* 1999; **159**: 1257–66.
- 61 Braun-Fahrlander C, Vuille JC, Sennhauser FH, et al. Respiratory health and long-term exposure to air pollutants in Swiss schoolchildren. *Am J Respir Crit Care Med* 1997; **155**: 1042–49.
- 62 Dockery DW, Cunningham J, Damokosh AI, et al. Health effects of acid aerosols on North American children: respiratory symptoms. *Environ Health Perspect* 1996; **104**: 500–05.
- 63 Raizenne M, Neas LM, Damokosh AI, et al. Health effects of acid aerosols on North American children: pulmonary function. *Environ Health Perspect* 1996; **104**: 506–14.
- 64 Spengler JD, Koutrakis P, Dockery DW, Raizenne M, Speizer FE. Health effects of acid aerosols on North American children: air pollution exposures. *Environ Health Perspect* 1996; **104**: 492–99.
- 65 Gauderman WJ, McConnell R, Gilliland F, et al. Association between air pollution and lung function growth in southern California children. *Am J Respir Crit Care Med* 2000; **162**: 1383–90.
- 66 Avol EL, Gauderman WJ, Tan SM, London SJ, Peters JM. Respiratory effects of relocating to areas of differing air pollution levels. *Am J Respir Crit Care Med* 2001; **164**: 2067–72.
- 67 Kaur B, Anderson HR, Austin J, et al. Prevalence of asthma symptoms, diagnosis, and treatment in 12–14 year old children across Great Britain (international study of asthma and allergies in childhood, ISAAC UK). *BMJ* 1998; **316**: 118–24.
- 68 The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. *Lancet* 1998; **351**: 1225–32.
- 69 McDonnell WF, Abbey DE, Nishino N, Lebowitz MD. Long-term ambient ozone concentration and the incidence of asthma in nonsmoking adults: the AHSMOG Study. *Environ Res* 1999; **80**: 110–21.
- 70 McConnell R, Berhane K, Gilliland F, et al. Asthma in exercising children exposed to ozone: a cohort study. *Lancet* 2002; **359**: 386–91.
- 71 Rahman I, MacNee W. Oxidative stress and regulation of glutathione in lung inflammation. *Eur Respir J* 2000; **16**: 534–54.
- 72 Bayram H, Sapsford RJ, Abdelaziz MM, Khair OA. Effect of ozone and nitrogen dioxide on the release of proinflammatory mediators from bronchial epithelial cells of nonatopic nonasthmatic subjects and atopic asthmatic patients in vitro. *J Allergy Clin Immunol* 2001; **107**: 287–94.
- 73 Mochitate K, Katagiri K, Miura T. Impairment of microbial killing and superoxide-producing activities of alveolar macrophages by a low level of ozone. *J Health Sci* 2001; **47**: 302–09.
- 74 Nichols BG, Woods JS, Lucht DL, Corral J, Koenig JQ. Effects of ozone exposure on nuclear factor-kappaB activation and tumor necrosis factor-alpha expression in human nasal epithelial cells. *Toxicol Sci* 2001; **60**: 356–62.
- 75 Bassett D, Elbon-Copp C, Otterbein S, et al. Inflammatory cell availability affects ozone-induced lung damage. *J Toxicol Environ Health A* 2001; **64**: 547–65.
- 76 Jorres R, Nowak D, Magnussen H. The effect of ozone exposure on allergen responsiveness in subjects with asthma or rhinitis. *Am J Respir Crit Care Med* 1996; **153**: 56–64.
- 77 Kleeberger SR, Levitt RC, Zhang LY, et al. Linkage analysis of susceptibility to ozone-induced lung inflammation in inbred mice. *Nat Genet* 1997; **17**: 475–78.
- 78 Cho HY, Zhang LY, Kleeberger SR. Ozone-induced lung inflammation and hyperreactivity are mediated via tumor necrosis factor-alpha receptors. *Am J Physiol Lung Cell Mol Physiol* 2001; **280**: L537–46.
- 79 Bergamaschi E, Palma G De, Mozzoni P, et al. Polymorphism of quinone-metabolizing enzymes and susceptibility to ozone-induced acute effects. *Am J Respir Crit Care Med* 2001; **163**: 1426–31.
- 80 Mudway IS, Krishna MT, Frew AJ, et al. Compromised concentrations of ascorbate in fluid lining the respiratory tract in human subjects after exposure to ozone. *Occup Environ Med* 1999; **56**: 473–81.
- 81 Samet JM, Hatch GE, Horstman D, et al. Effect of antioxidant supplementation on ozone-induced lung injury in human subjects. *Am J Respir Crit Care Med* 2001; **164**: 819–25.
- 82 Holz O, Jorres RA, Timm P, et al. Ozone-induced airway inflammatory changes differ between individuals and are reproducible. *Am J Respir Crit Care Med* 1999; **159**: 776–84.
- 83 Schelegle ES, Eldridge MW, Cross CE, Walby WF, Adams WC. Differential effects of airway anesthesia on ozone-induced pulmonary responses in human subjects. *Am J Respir Crit Care Med* 2001; **163**: 1121–27.
- 84 Jorres RA, Holtz O, Zachgo P, et al. The effect of repeated ozone exposures on inflammatory markers in bronchoalveolar lavage fluid and mucosal biopsies. *Am J Respir Crit Care Med* 2000; **161**: 1855–61.
- 85 Blomberg A, Krishna MT, Bocchino V, et al. The inflammatory effects of 2 ppm NO2 on the airways of healthy subjects. *Am J Respir Crit Care Med* 1997; **156**: 418–24.
- 86 Sandstrom T, Helleday R, Bjerner L, Stjernberg N. Effects of repeated exposure to 4 ppm nitrogen dioxide on bronchoalveolar lymphocyte subsets and macrophages in healthy men. *Eur Respir J* 1992; **5**: 1092–96.
- 87 Frampton MW, Smeglin AM, Roberts NJ Jr, Finkelstein JN, Morrow PE, Utell MJ. Nitrogen dioxide exposure in vivo and human alveolar macrophage inactivation of influenza virus in vitro. *Environ Res* 1989; **48**: 179–92.

- 88 Chauhan AJ, Krishna MT, Frew AJ, Holgate ST. Exposure to nitrogen dioxide (NO<sub>2</sub>) and respiratory disease risk. *Rev Environ Health* 1998; **13**: 73–90.
- 89 Tunnicliffe WS, Burge PS, Ayres JG. Effect of domestic concentrations of nitrogen dioxide on airway responses to inhaled allergen in asthmatic patients. *Lancet* 1994; **344**: 1733–36.
- 90 Fujii T, Hayashi S, Hogg JC, Vincent R, Van Eeden SF. Particulate matter induces cytokine expression in human bronchial epithelial cells. *Am J Respir Cell Mol Biol* 2001; **25**: 265–71.
- 91 Lundborg M, Johard U, Lastbom L, Gerde P, Camner P. Human alveolar macrophage phagocytic function is impaired by aggregates of ultrafine carbon particles. *Environ Res* 2001; **86**: 244–53.
- 92 Stearns RC, Paulauskis JD, Godleski JJ. Endocytosis of ultrafine particles by A549 cells. *Am J Respir Cell Mol Biol* 2001; **24**: 108–15.
- 93 Renwick LC, Donaldson K, Clouter A. Impairment of alveolar macrophage phagocytosis by ultrafine particles. *Toxicol Appl Pharm* 2001; **172**: 119–27.
- 94 Beck-Speier I, Dayal N, Karg E, et al. Agglomerates of ultrafine particles of elemental carbon and TiO<sub>2</sub> induce generation of lipid mediators in alveolar macrophages. *Environ Health Perspect* 2001; **109** (suppl 4): 613–18.
- 95 Dellinger B, Pryor WA, Cueto R, et al. Role of free radicals in the toxicity of airborne fine particulate matter. *Chem Res Toxicol* 2001; **14**: 1371–77.
- 96 Sun G, Crissman K, Norwood J, et al. Oxidative interactions of synthetic lung epithelial lining fluid with metal-containing particulate matter. *Am J Physiol Lung Cell Mol Physiol* 2001; **281**: L807–15.
- 97 Van Eeden SF, Tan WC, Suwa T, et al. Cytokines involved in the systemic inflammatory response induced by exposure to particulate matter air pollutants (PM<sub>10</sub>). *Am J Respir Crit Care Med* 2001; **164**: 826–30.
- 98 Salvi SS, Nordenhall C, Blomberg A, et al. Acute exposure to diesel exhaust increases IL-8 and GRO- $\alpha$  production in healthy human airways. *Am J Respir Crit Care Med* 2000; **161**: 550–57.
- 99 Salvi S, Blomberg A, Rudell B, et al. Acute inflammatory responses in the airways and peripheral blood after short-term exposure to diesel exhaust in healthy human volunteers. *Am J Respir Crit Care Med* 1999; **159**: 702–09.
- 100 Mukae H, Vincent R, Quinlan K, et al. The effect of repeated exposure to particulate air pollution (PM<sub>10</sub>) on the bone marrow. *Am J Respir Crit Care Med* 2001; **163**: 201–09.
- 101 Wu W, Samet JM, Ghio AJ, Devlin RB. Activation of the EGF receptor signaling pathway in airway epithelial cells exposed to Utah Valley PM. *Am J Physiol Lung Cell Mol Physiol* 2001; **281**: L483–89.
- 102 Puddicombe SM, Davies DE. The role of MAP kinases in intracellular signal transduction in bronchial epithelium. *Clin Exp Allergy* 2000; **30**: 7–11.
- 103 Seaton A, Soutar A, Crawford V, et al. Particulate air pollution and the blood. *Thorax* 1999; **54**: 1027–32.
- 104 Hashimoto K, Ishii Y, Uchida Y, et al. Exposure to diesel exhaust exacerbates allergen-induced airway responses in guinea pigs. *Am J Respir Crit Care Med* 2001; **164**: 1957–63.
- 105 Fujieda S, Diaz-Sanchez D, Saxon A. Combined nasal challenge with diesel exhaust particles and allergen induces in vivo IgE isotype switching. *Am J Respir Cell Mol Biol* 1998; **19**: 507–12.
- 106 Nel AE, Diaz-Sanchez D, Ng D, Hiura T, Saxon A. Enhancement of allergic inflammation by the interaction between diesel exhaust particles and the immune system. *J Allergy Clin Immunol* 1998; **102**: 539–54.
- 107 Brown DM, Wilson MR, MacNee W, Stone V, Donaldson K. Size-dependent proinflammatory effects of ultrafine polystyrene particles: a role for surface area and oxidative stress in the enhanced activity of ultrafines. *Toxicol Appl Pharm* 2001; **175**: 191–99.
- 108 Cakmak S, Burnett RT, Krewski D. Methods for detecting and estimating population threshold concentrations for air pollution-related mortality with exposure measurement error. *Risk Anal* 1999; **19**: 487–96.
- 109 Daniels MJ, Dominici F, Samet JM, Zeger SL. Estimating particulate matter-mortality dose-response curves and threshold levels: an analysis of daily time-series for the 20 largest US cities. *Am J Epidemiol* 2000; **152**: 397–406.
- 110 Brunekreef B, Dockery DW, Krzyzanowski M. Epidemiologic studies on short-term effects of low levels of major ambient air pollution components. *Environ Health Perspect* 1995; **103** (suppl 2): 3–13.
- 111 McMichael AJ, Anderson HR, Brunekreef B, Cohen AJ. Inappropriate use of daily mortality analyses to estimate longer-term mortality effects of air pollution. *Int J Epidemiol* 1998; **27**: 450–53.
- 112 Schwartz J. Harvesting and long term exposure effects in the relation between air pollution and mortality. *Am J Epidemiol* 2000; **151**: 440–48.
- 113 Schwartz J. Is there harvesting in the association of airborne particles with daily deaths and hospital admissions? *Epidemiology* 2001; **12**: 55–61.
- 114 Zanobetti A, Schwartz J, Samoli E, et al. The temporal pattern of mortality responses to air pollution: a multicity assessment of mortality displacement. *Epidemiology* 2002; **13**: 87–93.
- 115 Zeger SL, Dominici F, Samet J. Harvesting-resistant estimates of air pollution effects on mortality. *Epidemiology* 1999; **10**: 171–75.
- 116 Schwartz J. What are people dying of on high air pollution days? *Environ Res* 1994; **64**: 26–35.
- 117 Janssen NA, Hartog JJ de, Hoek G, et al. Personal exposure to fine particulate matter in elderly subjects: relation between personal, indoor, and outdoor concentrations. *J Air Waste Manag Assoc* 2000; **50**: 1133–43.
- 118 Sarnat JA, Schwartz J, Catalano PJ, Suh HH. Gaseous pollutants in particulate matter epidemiology: confounders or surrogates? *Environ Health Perspect* 2001; **109**: 1053–61.
- 119 Rijnders E, Janssen NA, Vliet PH van, Brunekreef B. Personal and outdoor nitrogen dioxide concentrations in relation to degree of urbanization and traffic density. *Environ Health Perspect* 2001; **109** (suppl 3): 411–17.
- 120 Oglesby L, Kunzli N, Roosli M, et al. Validity of ambient levels of fine particles as surrogate for personal exposure to outdoor air pollution—results of the European EXPOLIS-EAS Study (Swiss Center Basel). *J Air Waste Manag Assoc* 2000; **50**: 1251–61.
- 121 Hoek G, Fischer P, Van Den Brandt P, Goldbohm S, Brunekreef B. Estimation of long-term average exposure to outdoor air pollution for a cohort study on mortality. *J Expo Anal Environ Epidemiol* 2001; **11**: 459–69.
- 122 Seaton A, MacNee W, Donaldson K, Godden D. Particulate air pollution and acute health effects. *Lancet* 1995; **345**: 176–78.
- 123 Peters A, Wichmann HE, Tuch T, Heinrich J, Heyder J. Respiratory effects are associated with the number of ultrafine particles. *Am J Respir Crit Care Med* 1997; **155**: 1376–83.
- 124 Penttinen P, Timonen KL, Tiittanen P, et al. Number concentration and size of particles in urban air: effects on spirometric lung function in adult asthmatic subjects. *Environ Health Perspect* 2001; **109**: 319–23.
- 125 van Vliet P, Knape M, de Hartog J, et al. Motor vehicle exhaust and chronic respiratory symptoms in children living near freeways. *Environ Res* 1997; **74**: 122–32.
- 126 Janssen NA, Schwartz J, Zanobetti A, Suh HH. Air conditioning and source-specific particles as modifiers of the effect of PM<sub>10</sub> on hospital admissions for heart and lung disease. *Environ Health Perspect* 2002; **110**: 43–49.
- 127 Roemer WH, van Wijnen JH. Daily mortality and air pollution along busy streets in Amsterdam, 1987–1998. *Epidemiology* 2001; **12**: 649–53.
- 128 Laden F, Neas LM, Dockery DW, Schwartz J. Association of fine particulate matter from different sources with daily mortality in six US cities. *Environ Health Perspect* 2000; **108**: 941–47.
- 129 Ghio AJ, Devlin RB. Inflammatory lung injury after bronchial instillation of air pollution particles. *Am J Respir Crit Care Med* 2001; **164**: 704–08.
- 130 Schwartz J, Dockery DW, Neas LM. Is daily mortality associated specifically with fine particles? *J Air Waste Manag Assoc* 1996; **46**: 927–39.
- 131 Schwartz J, Neas LM. Fine particles are more strongly associated than coarse particles with acute respiratory health effects in schoolchildren. *Epidemiology* 2000; **11**: 6–10.
- 132 McDonnell WF, Nishino-Ishikawa N, Petersen FF, Chen LH, Abbey DE. Relationships of mortality with the fine and coarse fractions of long-term ambient PM<sub>10</sub> concentrations in nonsmokers. *J Expo Anal Environ Epidemiol* 2000; **10**: 427–36.
- 133 Smith RL, Spitzner D, Kim Y, Fuentes M. Threshold dependence of mortality effects for fine and coarse particles in Phoenix, Arizona. *J Air Waste Manag Assoc* 2000; **50**: 1367–79.
- 134 Anderson HR, Bremner SA, Atkinson RW, Harrison RM, Walters S. Particulate matter and daily mortality and hospital admissions in the west Midlands conurbation of the United Kingdom: associations with fine and coarse particles, black smoke and sulphate. *Occup Environ Med* 2001; **58**: 504–10.
- 135 Pope CA 3rd. Respiratory disease associated with community air pollution and a steel mill. *Utah Val Am J Public Health* 1989; **79**: 623–28.
- 136 Pope CA 3rd. Particulate pollution and health: a review of the Utah valley experience. *J Expo Anal Environ Epidemiol* 1996; **6**: 23–34.
- 137 Friedman MS, Powell KE, Hutwagner L, Graham LM, Teague WG. Impact of changes in transportation and commuting behaviors during the 1996 Summer Olympic Games in Atlanta on air quality and childhood asthma. *JAMA* 2001; **285**: 897–905.
- 138 Heinrich J, Hoelscher B, Wichmann HE. Decline of ambient air

- pollution and respiratory symptoms in children. *Am J Respir Crit Care Med* 2000; **161**: 1930–36.
- 139 Schwartz J, Dockery DW. Increased mortality in Philadelphia associated with daily air pollution concentrations. *Am Rev Respir Dis* 1992; **145**: 600–04.
- 140 Kelsall JE, Samet JM, Zeger SL, Xu J. Air pollution and mortality in Philadelphia, 1974–1988. *Am J Epidemiol* 1997; **146**: 750–62.
- 141 Krewski D, Burnett R, Goldberg MS, et al. Reanalysis of the Harvard six cities study and the American Cancer Society study of particulate air pollution and mortality: Health Effects Institute special report. Boston: Health Effects Institute, 2000.
- 142 Samet J, Zeger S, Kelsall J, Xu J, Kalkstein L. Does weather confound or modify the association of particulate air pollution with mortality? An analysis of the Philadelphia data, 1973–1980. *Environ Res* 1998; **77**: 9–19.
- 143 Pope CA 3rd, Kalkstein LS. Synoptic weather modeling and estimates of the exposure–response relationship between daily mortality and particulate air pollution. *Environ Health Perspect* 1996; **104**: 414–20.
- 144 Burnett R, Ma R, Jerrett M, et al. The spatial association between community air pollution and mortality: a new method of analyzing correlated geographic cohort data. *Environ Health Perspect* 2001; **109** (suppl 3): 375–80.
- 145 Greenbaum DS, Bachmann JD, Krewski D, et al. Particulate air pollution standards and morbidity and mortality: case study. *Am J Epidemiol* 2001; **154**: S78–90.
- 146 Kunzli N, Kaiser R, Medina S, et al. Public-health impact of outdoor and traffic-related air pollution: a European assessment. *Lancet* 2000; **356**: 795–801.
- 147 Hollander AE de, Melse JM, Lebrete E, Kramers PG. An aggregate public health indicator to represent the impact of multiple environmental exposures. *Epidemiology* 1999; **10**: 606–17.

## Uses of error

### Uncertainty

David A Grimes

Everything pointed to the same diagnosis: an ectopic pregnancy. Nonetheless, I was nervous making the diagnosis by myself. I was a very junior house officer in obstetrics and gynaecology, alone on call at a municipal hospital. My patient had an abnormal last menses about 8 weeks ago, no recent use of contraception, and vague pelvic pain. Her pulse and blood pressure were normal without orthostatic changes. She looked uncomfortable but not in acute distress. Her abdomen was tender, and the pelvic examination caused her discomfort, more on one side than the other. I could feel no adnexal mass, but my clinical experience was very limited. Her haemoglobin was around 110 g/L; not unusual for our indigent patients.

From my readings and observations as a medical student, I knew just what to do to confirm my diagnosis: a culdocentesis. At this time, sensitive pregnancy tests were not available, diagnostic ultrasound was in its infancy, and laparoscopy had just reached the USA and was not available by night at my hospital. After obtaining informed consent, I introduced a speculum and inserted a 20-gauge needle through her vaginal vault into her rectouterine pouch. To my delight, when I withdrew the plunger, the syringe promptly filled with dark blood. To my considerable dismay, however, the blood promptly clotted in the syringe. This was not supposed to happen: the hallmark of an ectopic pregnancy was non-clotting blood on culdocentesis. I had never read, seen, or heard of blood clotting in this setting.

Alone with this diagnostic dilemma, I excused myself from the patient and her family and slipped off discreetly to the hospital library. My memory had been correct: every gynaecology textbook on the shelf concurred that non-clotting blood on culdocentesis indicated an ectopic pregnancy. None, however, mentioned my current quandary: intraperitoneal blood that clotted. Given this uncertainty, I did not feel comfortable mobilising an operating theatre (which required calling in staff from home) to perform a laparotomy. Nor did I feel comfortable sending my patient home. I admitted her to the hospital, where I watched and fretted over her all night.

By morning, her pain was unchanged. Her repeat haemoglobin, however, had plummeted, and my error was apparent. We rushed her to the operating theatre. At laparotomy, she had litres of blood in her abdomen and required transfusion of several units of blood as a consequence of my timidity. I had erred in trusting the textbooks instead of my inexperienced clinical hunch. I later learned from published articles that a few such patients have blood that will not clot. Some authors call this “non-conclusive” evidence of an ectopic pregnancy.

I learned an important lesson that night about the limitations of textbooks and authorities. As Antman and colleagues showed several decades later, textbooks and their authors are dangerously obsolete. But Mark Twain may have put it best a century ago: “Be careful about reading health books. You may die of a misprint.” The same holds true for omissions.

Family Health International, Research Triangle Park, NC 27709, USA (D A Grimes MD)