



Total laryngectomy is generally done for advanced cancers of the larynx and hypopharynx, recurrence following (chemo)radiation, and occasionally for intractable aspiration and advanced thyroid cancer invading the larynx.

Although it is an excellent oncologic procedure and secures good swallowing without aspiration, it has disadvantages such as having a permanent tracheostomy; that verbal communication is dependent on oesophageal speech, and/or tracheo-oesophageal fistula speech or an electrolarynx; hyposmia; and the psychological and financial/ employment implications. Even in the best centers, about 20% of patients do not acquire useful verbal communication.

Prelaryngectomy decision making

The surgeon needs to consider the following issues before embarking on a laryngectomy.

What will be the tumour resection lines?

As the initial incisions into the pharynx are done from externally without having the tumour in view, the surgeon must carefully assess the valleculae, base of tongue and the pyriform fossae for tumour involvement, so as to avoid cutting into tumour when entering the pharynx. Involvement of the base of tongue may also prompt the surgeon to opt for a retrograde laryngectomy (commencing the laryngectomy at the tracheostomal end of the specimen). In the absence of CT or MRI imaging, one can palpate and assess tumour involvement of the preepiglottic space and base of tongue under general anaesthesia by placing one index finger in the valleculae, and the other on the skin of the neck just above the hyoid bone. The

fingers should normally virtually meet, unless there is tumour in the preepiglottic space or vallecula or base of tongue.

Is thyroidectomy required? Both hypothyroidism and hypoparathyroidism are common sequelae of total laryngectomy, particularly following postoperative radiation therapy, and may be difficult to manage in a developing world setting. Twenty-five percent of laryngectomy patients become hypothyroid following hemithyroidectomy; and 75% if postoperative radiation is added. However both thyroid lobes may be preserved unless Level 6 nodes need to be resected with subglottic and pyriform fossa carcinoma, or when there is intraoperative or radiological evidence of direct tumour extension to involve the thyroid gland.

Will a pectoralis major flap be required?

A capacious pharynx is essential for good swallowing and fistula speech. Should tumour involve the hypopharynx, especially when it extends distally towards the cricopharyngeus, then the expertise has to be available to possibly augment the pharyngeal repair with a pectoralis major flap. Pectoralis major muscle flaps are also frequently used to overlay the pharyngeal repair with salvage laryngectomy to reduce the fistula rate.

Is elective neck dissection required? With advanced laryngeal squamous cell carcinoma requiring laryngectomy, elective lateral neck dissection (Levels 2-4), either ipsilateral (glottic carcinoma) or bilateral (supraglottic, medial wall of pyriform fossa, bilateral glottic carcinoma) is recommended, with conversion to modified neck dissection should cervical metastases be found intraoperatively. Level 6 is included in subglottic and

pyriform fossa carcinoma to clear the paratracheal nodes.

Is the patient suitable for tracheo-oesophageal speech? This decision is based on assessment of cognitive function, motivation, financial ability to pay for replacement speech prostheses, and proximity to speech services.

Are there synchronous primaries or distant metastases? Total laryngectomy has significant morbidity, and should only be done if panendoscopy and CXR/CT chest exclude metastases or 2nd primaries.

Anaesthesia

Intubation: The operation is done under general anaesthesia. The ENT surgeon must be present to assist with a possibly difficult intubation. If a difficult intubation is anticipated, then either do an awake tracheostomy, or infiltrate skin and trachea with local anaesthesia/vasoconstrictor, in preparation for a possible emergency tracheostomy.

Preoperative tracheotomy: Tracheotomy may have been required for airway obstruction. It is not an independent indication for postoperative radiation therapy unless tumour was entered at the time of tracheotomy. If a tracheotomy has already been done, then ask the anaesthetist to reintubate through the larynx with an orotracheal tube once the patient has been anaesthetised as this facilitates dissection in the lower neck and speeds up the surgery.

Perioperative antibiotics: Commence perioperative antibiotics before putting knife to skin, and continue for 24 hrs

Surgical anatomy

Figures 1 & 2 illustrate all the muscles that will be divided during laryngectomy.




Figure 1: Supra- and infrahyoid muscles




Figure 2: Middle and inferior pharyngeal constrictors

Surgical steps

Positioning: Hyperextend the neck

Incisions for apron flap (Figures 3a, b)

The horizontal limb of the flap is placed approximately 2cms above the sternal notch. An ellipse of skin around a pre-existing tracheostomy is included with the resection. With a simple laryngectomy the vertical incision are placed along the

anterior borders of sternocleidomastoid muscles. For a laryngectomy with neck dissection(s), either a wider flap overlying the sternocleidomastoid muscles is made (*Figure 3a*), or a narrow flap with inferolateral extensions is made (*Figure 3b*). The latter has the disadvantage of a trifurcation which is more prone to wound breakdown and exposure of the major cervical vessels.



Figure 3a: Wide apron flap to accommodate neck dissections




Figure 3b: Narrow apron flap for laryngectomy, with lateral extensions for neck dissections

Flap elevation (*Figure 4*)

- Cut through the superficial layer of investing fascia and platysma muscles. The platysma is often absent in

midline. Take care not to injure the external and anterior jugular veins.

- Elevate the apron flap in a subplatysmal plane, remaining superficial to the external and anterior jugular veins.
- Dissect the flap superiorly up to approximately 2cms above the body of the hyoid bone.




Figure 4: Elevated apron flap and incisions through investing layer of cervical along anterior borders of sternocleidomastoid muscles

Freeing up the larynx

Free up one side of the larynx at a time. Stand on the side of neck that is being dissected.

- Ligate and transect the anterior jugular veins suprasternally and above the hyoid.
- Incise the investing layer of cervical fascia along the anterior border of the sternocleidomastoid muscle (*Figure 4*).
- Retract the sternocleidomastoid muscle laterally
- Identify the sternohyoid and omohyoid muscles
- Transect the omohyoid muscle medial to where it crosses the internal jugular vein (*Figure 5*)




Figure 5: Transect omohyoid along yellow line

- Identify the dissection plane between carotid sheath and larynx and thyroid gland and open this plane with sharp and blunt dissection with a finger to expose prevertebral fascia (Figure 6)




Figure 6: Transect sternohyoid muscle to expose sternohyoid muscle

- Transect the sternohyoid muscle with electrocautery wherever convenient (Figure 6)
- Identify the sternohyoid muscle, and carefully divide it below larynx (Figure 6). It is a broad, thin muscle, so take special care not to injure the thyroid gland and its rich vasculature which is immediately deep to muscle

- Carefully elevate and reflect the superior cut end of the sternohyoid muscle from the thyroid gland using electrocautery dissection (Figure 7)




Figure 7: Transect & elevate sternohyoid to expose thyroid gland




Figure 8: Divided sternohyoid retracted to expose thyroid. Line indicates course of dissection of thyroid gland and along midline of trachea

- Divide the thyroid isthmus with electrocautery
- Divide and strip the tissues overlying the cervical trachea anteriorly in the

midline to avoid injuring the inferior thyroid veins

- Carefully reflect the thyroid lobe off the trachea, cricoid and inferior constrictor with electrocautery (*Figure 9*) while inspecting for and excluding direct laryngeal tumour extension to the thyroid gland




Figure 9: Thyroid gland has been mobilised from larynx and trachea

- Identify and transect the recurrent laryngeal nerve (*Figure 10*)
- Identify the tracheo-oesophageal groove and oesophagus (*Figure 10*)




Figure 10: Identify oesophagus, and divide recurrent laryngeal nerve

- Identify and divide the superior laryngeal branch of superior thyroid

artery, and reflect and preserve the superior thyroid pedicle from the larynx (*Figure 11*)

- Identify and divide the superior laryngeal nerve




Figure 11: Identify and divide superior laryngeal branch of superior thyroid artery

- Rotate the larynx to the contralateral side, and identify the posterior border of the thyroid ala (*Figure 12*)




Figure 12: Rotate the larynx with a finger placed behind the thyroid ala

- Divide the inferior pharyngeal constrictor muscle and thyroid perichondrium with electrocautery at,

or just anterior to the posterior border of the thyroid ala (*Figure 13*)




Figure 13: Divided inferior pharyngeal constrictor and thyroid perichondrium

- Strip the lateral wall of the pyriform fossa off the medial aspect of the thyroid ala in a subperichondrial plane with a swab/sponge held over a fingertip, or with a Freer's elevator, **only** on the side of the larynx opposite to the cancer (*Figure 14*). On the side of the cancer, this step is omitted to ensure adequate resection margins.




Figure 14: Pyriform fossa mucosa stripped from thyroid lamina

The surgeon then crosses to the opposite side of the patient, and repeats the above operative steps.

Suprahyoid dissection

The following description applies to laryngeal cancer not involving the preepiglottic space, vallecula or the base of tongue. When tumour does involve vallecula, pre-epiglottic space and/or base of tongue, then the pharynx is entered via the opposite pyriform fossa or a retrograde laryngectomy is done, commencing the dissection inferiorly at tracheostomy.

- Identify the body of the hyoid bone. Remember that the hypoglossal nerves and lingual arteries lie deep to the greater cornua/horns of the hyoid bone
- Divide the suprahyoid muscles with electrocautery along the superior border of the body of the hyoid bone (*Figure 15*)




Figure 15: Transection of suprahyoid muscles from hyoid body

- Initially do not dissect lateral to the lesser cornua, as the hypoglossal nerves and the lingual arteries are located deep to the greater cornua of the hyoid bone
- Release the digastric tendon and stylohyoid ligament and muscle from the lesser cornu of the hyoid. The hyoid then become more mobile and can be displaced inferiorly, away from the hypoglossal nerves
- Rotate the hyoid bone to the contralateral side, and identify the position of the greater cornu/horn of the hyoid bone (*Figure 16*)




Figure 16: Identify greater cornu

- The hyoglossus and middle constrictor muscles are next released from the greater cornu with diathermy
- Divide the soft tissue on the medial aspect of the tips of the greater cornua of the hyoid with scissors to isolate the greater cornua of the hyoid bilaterally (*Figure 17*). Hug the inner aspect of the greater cornua to avoid the hypoglossal nerves. If a neck dissection has been done, the hypoglossal nerves will already be visible




Figure 17: Releasing greater cornu

- Dissect transversely with diathermy along the superior margin of the body of the hyoid bone, and along the superior margin of the pre-epiglottic space. Identify the hyoepiglottic ligament in the midline. Dissect along the hyoepiglottic ligament and strip the vallecula mucosa from the anterior surface of the epiglottis (*Figure 18*)




Figure 18: Suprahyoid approach to valleculae

- Enter the pharynx by incising the mucosa along the superior margin of the epiglottis (*Figure 19*)




Figure 19: Entering vallecula

Tracheostomy

- A tracheostomy is done at this stage so as to mobilise the larynx and to facilitate the laryngeal resection
- Ask the anaesthetist to preoxygenate the patient
- Incise the trachea transversely between the 3rd/4th/5th tracheal rings or below a preoperative tracheostomy. With a small trachea, incise the lateral tracheal walls in a superolateral direction to bevel and enlarge the tracheostoma. Place a few 3-0 vicryl half-mattress sutures between the anterior wall of the transected trachea and the skin to approximate mucosa to skin
- Puncture and deflate the cuff of the endotracheal tube, and cut the tube in the pharynx, and remove the distal end of the tube through the pharyngotomy
- Insert a flexible endotracheal tube e.g. armoured tube into the tracheostoma. Avoid inserting the tube too deeply as the carina is quite close to the tracheostoma. Fix the tube to the chest wall or drapes with a temporary suture so that it does not become displaced, attach the sterile anaesthesia tubing and resume ventilation

Laryngeal resection

- Inspect the subglottis through the tracheostoma to ensure that the tracheal resection margin is adequate
- Move to the head of the operating table
- Retract the epiglottis and the larynx anteriorly through the pharyngotomy, and inspect the larynx and the tumour
- Commence laryngeal resection contralateral to the tumour using curved scissors with points located anteriorly/upwards so as to avoid inadvertently resecting too much pharyngeal mucosa
- Cut along the lateral border of the epiglottis on the less involved side, to expose the hypopharynx
- Repeat this on the side of tumour, with at least a 1cm mucosal margin around the tumour
- On the less involved side, cut through the lateral wall of the pyriform fossa and hug the arytenoids and cricoid to preserve pyriform sinus mucosa (Figure 20). The superior laryngeal neurovascular pedicle will be transected if not previously addressed
- Repeat on the tumour side




Figure 20: Resect the larynx preserving maximum amount of pharyngeal mucosa

- Join the left and right pyriform incisions by tunnelling below and cutting the postcricoid mucosa transversely (Figure 21)




Figure 21: Transverse postcricoid cut

- Separate the posterior wall of the larynx (cricoid, tracheal membrane) from the anterior wall of the oesophagus by dissecting with a scalpel along the avascular plane between that exists between oesophagus and trachea/cricoid (*Figure 22*). Take care to stop just short of the tracheostoma.




Figure 22: Dissecting in the avascular plane between oesophagus and trachea

- Transect the posterior wall of the trachea, and remove the larynx (*Figure 23*)
- Inspect the laryngectomy specimen for adequacy of resection margins, and resect additional tissue if indicated




Figure 23: Transect trachea and remove larynx

Pharyngo-oesophageal myotomy

- Optimising speech and swallowing requires a capacious and floppy pharynx
- Always perform a pharyngo-oesophageal myotomy to prevent hypertonicity of the pharyngo-oesophageal segment
- Insert an index finger into the oesophagus
- With a sharp scalpel, divide all the muscle fibres down to the submucosa, and distally to the level of the tracheostoma (*Figure 24*). The myotomy may be done in the midline or to the side.



Figure 24: Cricopharyngeal myotomy

Tracheo-oesophageal fistula

- Tracheo-oesophageal speech is the best form of alaryngeal communication
- A tracheo-oesophageal fistula is created before closing the pharynx
- Pass a curved artery forceps through the pharyngeal defect and along the oesophagus, and tent up the anterior wall of oesophagus/posterior tracheal wall 5-10mm below the superior margin of the tracheostoma. Placing the fistula too low makes changing the prosthesis difficult
- Cut down onto the artery tip with a scalpel, and pass the tip of the artery forceps through the fistula into the tracheal lumen
- Hold the tip of a 14 gauge Foley urinary catheter with the artery forceps, and pull the catheter through the fistula into the oesophagus and pass it through the pharyngeal defect (*Figure 25*). Then advance the catheter down the oesophagus. Avoid accidental displacement of the catheter by injecting 5ml water into the bulb and by fixing the catheter to the skin with a suture




Figure 25: Creation of tracheo-oesophageal fistula

- The catheter acts a stent to allow the fistula to mature in preparation for fitting of a tracheo-oesophageal

prosthesis, and is initially used for stomagastric feeding

- An alternative method is to insert a speech prosthesis *ab initio*, and to feed the patient via a nasogastric tube, or a catheter passed through the speech prosthesis ([Postlaryngectomy vocal and pulmonary rehabilitation](#))
- Divide the sternal heads of the sternomastoid muscles to create a flattened peristomal contour and to facilitate digital stomal occlusion (*Figure 26*).




Figure 26: Division of sternal heads of sternomastoid to flatten peristomal area

Pharyngeal closure

- At least 2.5cm transverse diameter of residual pharyngeal mucosa is required for primary pharyngeal closure. The teaching that the minimum pharynx required is that which may be closed over a nasogastric tube is incorrect, as the neopharynx is then too narrow for adequate swallowing and voicing
- A horizontal/transverse closure is preferred as it maximises the capacity of the pharynx (*Figures 27*). Only if there is undue tension on the suture line, then do T-shaped closure, keeping the vertical limb as short as possible.




Figure 27: Pharynx well suited to a transverse closure

- Take care not to injure the lingual arteries when suturing the pharynx, as injury to the arteries may lead to necrosis of the tongue
- A 3-layered pharyngeal closure is suggested
 - 1st layer: 3-0 vicryl running modified Connell or true Connell technique (Invert mucosa)




Figure 28: Completed 1st layer of transverse closure of pharynx

- 2nd layer: 3-0 vicryl running suture of submucosa and muscle

- 3rd layer: Approximate inferior constrictors and suture constrictors to suprahyoid muscles with interrupted 3-0 vicryl

Final steps

- Ask the anaesthetist to do a Valsalva manoeuvre to exclude bleeding and chyle leaks
- If there is excessive, lax suprastomal skin that may occlude the tracheostomy when the patient flexes the neck, then trim a crescent of skin from the inferior edge of the apron flap suprastomally
- Suture the skin to the edges of the tracheostomy with half-mattress interrupted 3-0 vicryl sutures
- Seal the trifurcation at the lateral edge of the stoma with a suture as indicated below (Figure 29)




Figure 29: Suture technique to seal trifurcation between skin and side of tracheostoma

- Insert a 1/4" suction drain
- Irrigate neck with sterile water
- Reapproximate the platysma with 3-0 vicryl running sutures
- Close the skin with a running nylon suture or with skin staples
- Suction blood from trachea
- Insert a cuffed tracheostomy tube, and suture it to skin

Postoperative care

- Antibiotics x 24 hours
- Chest physiotherapy
- Remove suction drains when <50mls drainage per 24hrs (*See references*)
- Day 1: Mobilise to chair, remove urinary catheter
- Day 2: Commence oral feeding. Early oral feeding is safe, and does not cause pharyngocutaneous fistulae (*See references*)

Early feeding protocol	
Day 1	Intravenous general maintenance solution & stomagastric feeds
Day 2	Clear oral fluids & stomagastric feeds
Day 3	Free oral fluids
Day 4	Soft diet
Day 5	Normal diet

- Day 7: Remove sutures
- Day 10: Insert speaking valve
- Cover the stoma with a bib (*Figures 30, 31*)



Figure 30: Stoma and Foley catheter feeding tube one week following surgery



Figure 31: Bib in place

Pharyngeal reconstruction

Following resection of large pyriform fossa tumours (*Figure 32*), or tumours that extend close the cricopharyngeus, or involve the postcricoid area, only a narrow strip of mucosa may remain to reconstruct the neopharynx. If the residual pharyngeal mucosal is <2.5cms in width, then additional tissue is required to avoid pharyngeal stenosis, dysphagia and poor speech (*Figure 33*).

Reconstructive options include pectoralis major and latissimus dorsi flaps, or microvascular free tissue transfer flaps (radial forearm, anterolateral thigh). All these flaps can be used to augment the pharyngeal repair, or when the pharynx has been completely resected, may be tubed to entirely replace the pharynx (*Figures 34 – 37*). A contrast swallow X-ray is done on about day 7 to exclude an anastomotic leak before commencing oral feeding.




Figure 32: Large carcinoma of hypopharynx that will require pharyngeal reconstruction




Figure 33: Insufficient pharyngeal mucosa for primary closure of pharynx




Figure 34: Pectoralis major augmentation of pharynx




Figure 35: Tubed pectoralis major flap




Figure 36: Tubed free anterolateral thigh flap




Figure 37: Free jejunal flap

Author & Editor

Johan Fagan MBChB, FCORL, MMed
Professor and Chairman
Division of Otolaryngology
University of Cape Town
Cape Town
South Africa
johannes.fagan@uct.ac.za



The Open Access Atlas of Otolaryngology, Head & Neck Operative Surgery by [Johan Fagan \(Editor\)](#) johannes.fagan@uct.ac.za is licensed under a [Creative Commons Attribution - Non-Commercial 3.0 Unported License](#)

Useful references

- [Aswani J, Thandar MA, Otiti J, Fagan JJ. Early oral feeding following total laryngectomy. *J Laryngol Otol.* 2009; 123:333-338](#)
- A practical guide to post-laryngectomy vocal and pulmonary rehabilitation - Fourth Edition: [Postlaryngectomy vocal and pulmonary rehabilitation](#)
- [Fagan JJ, Lentin R, Oyarzabal MF, S Iaacs, Sellars SL. Tracheo-oesophageal speech in a Developing World Community. *Arch Otolaryngol* 2002, 128\(1\): 50-53](#)
- Fagan JJ, Kaye PV. Management of the thyroid gland with laryngectomy for cT3 glottic carcinomas. *Clin Otolaryngol*, 1997; 22: 7-12
- Harris T, Doolarkhan Z, Fagan JJ. Timing of removal of neck drains with head and neck surgery. *Ear Nose Throat J.* 2011 Apr; 90(4):186-9.